

every three hours, washing it down with three table-spoonsful of some febrifuge mixture.*

CHAP. XIX.

OPHTHALMIA, OR INFLAMMATION OF THE EYES

THIS disease may be occasioned by external injuries; as blows, burns, bruises, and the like. It may likewise proceed from dust, quick-lime, or other substances, getting into the eyes. It is often caused by the stoppage of customary evacuations; as the healing of old sores, drying up of issues, the suppressing of gentle morning sweats, or of the sweating of the feet, &c. Long exposure to the night air, especially in cold northerly winds, or whatever suddenly checks the perspiration, especially after the body has been much heated, is very apt to cause an inflammation of the eyes. Viewing snow, or other white bodies, for a long time, or looking steadfastly at the sun, a clear fire, or any bright object, will likewise occasion this malady. A sudden transition from darkness to very bright light will often have the same effect.

Nothing more certainly occasions an inflammation of the eyes than night-watching, especially reading or writing by candle-light. Drinking spirituous liquors, and excess of venery, are likewise very hurtful to the eyes. The acrid fumes of metals, and of several kinds of fuel, are also pernicious. Sometimes an inflammation of the eyes proceeds from a venereal taint, and often from a scrophulous or gouty habit. It may likewise be occasioned by hairs in the eye-lids turning inwards, and hurting the eyes. Sometimes the disease is epidemic, especially after wet seasons; and I have frequently known it prove infectious, particularly to those who lived in the same house with the patient. It may be occasioned by moist air, or living in low damp houses, especially in persons who are not accustomed to such situations. In children it often proceeds from imprudently drying up of scabbed heads, a running behind the ears, or any other discharge of that kind. Inflammations of the eyes often succeed the small-pox or measles, especially in children of a scrophulous habit.

SYMPTOMS.—An inflammation of the eyes is attended with acute pain, heat, redness, and swelling. The patient is not able to bear the light, and sometimes he feels a pricking pain, as if his eyes were pierced with a thorn. Sometimes he imagines his eyes are full of motes, or thinks he sees flies dancing before him. The eyes are filled with a scalding rheum, which rushes forth in great quantities, whenever the patient attempts to look up. The pulse

* Take Lemon Juice, $1\frac{1}{2}$ oz.
Subcarbonate of Ammonia $\frac{1}{2}$ dr.
Mint Water, 1 oz.
Pure Water, 4 oz.

Nitrate of Potash, 1 dr.
Syrup of Roses, 2 dra.
Make a mixture. To be taken as above

is generally quick and hard, with some degree of fever. When the disease is violent, the neighbouring parts swell, and there is a throbbing or pulsation in the temporal arteries, &c.

A slight inflammation of the eyes, especially from an external cause, is easily cured; but when the disease is violent, and continues long, it often leaves specks upon the eyes, or dimness of sight, and sometimes total blindness.

If the patient be seized with a looseness, it has a good effect; and when the inflammation passes from one eye to another, as it were by infection, it is no unfavourable symptom. But when the disease is accompanied with a violent pain in the head, and continues long, the patient is in danger of losing his sight.

REGIMEN.—The diet, unless in scrophulous cases, can hardly be too spare, especially at the beginning. The patient must abstain from every thing of a heating nature. His food should consist chiefly of mild vegetables, weak broths, and gruels. His drink may be barley-water, balm-tea, common whey, and such like.

The patient's chamber must be darkened, or his eyes shaded by a cover, so as to exclude the light, but not to press upon the eyes.* He should not look at a candle, the fire, or any luminous object; and ought to avoid all smoke, as the fumes of tobacco, or any thing that may cause coughing, sneezing, or vomiting. He should be kept quiet, avoiding all violent efforts, either of body or mind, and encouraging sleep as much as possible.

MEDICAL TREATMENT.—This is one of those diseases wherein great hurt is often done by external applications. Almost every person pretends to be possessed of a remedy for the cure of sore eyes. These remedies generally consist of eye-waters and ointments, with other external applications, which do mischief twenty times for once they do good. People ought, therefore, to be very cautious how they use such things, as even the pressure upon the eyes often increases the malady.

Bleeding, in a violent inflammation of the eyes, is always necessary. This should be performed as near the part affected as possible. An adult may lose ten or twelve ounces of blood from the jugular vein, and the operation may be repeated according to the urgency of the symptoms. If it should not be convenient to bleed in the neck, the same quantity may be let from the arm, or any other part of the body.

Leeches are often applied to the temples, or under the eyes, with good effect. The wounds must be suffered to bleed for some hours, and if the bleeding stop soon, it may be promoted by the application of cloths dipt in warm water. In obstinate cases, it will be necessary to repeat this operation several times.

Opening and diluting medicines are by no means to be neglected. The patient may take a small dose of Epsom salts, and cream

* The best kind of shade for tender eyes is formed by extending green gauze on wire properly constructed. By this contrivance the access of too much light is effectually impeded, while there is no interruption to the free access of the air, so that the eyes are not heated by this, as by the common shades of silk, or spectacles of green glass. A. P. B

of tartar, every second or third day, or a decoction of tamarinds with senna. If these be not agreeable, small doses of rhubarb and nitre, a little of the lenitive electuary, or any other mild purgative, will answer the same end. The patient at the same time must drink freely of water-gruel, tea, whey, or any other weak diluting liquor. He ought likewise to take, at bed-time, a large draught of very weak wine-whey, in order to promote perspiration. His feet and legs must frequently be bathed in lukewarm water, and his head shaved twice or thrice a-week, and afterwards washed in cold water. This has often a remarkably good effect.

If the inflammation does not yield to these evacuations, blisters must be applied to the temples, behind the ears, or upon the neck, and kept open for some time by the mild blistering-ointment, or they may be frequently renewed, which is often preferable. I have seldom known these, if long enough kept open, fail to remove the most obstinate inflammation of the eyes; but for this purpose it is often necessary to continue the discharge for several weeks.

To abate the inflammation, it is customary to have recourse to the frequent application of some cooling and astringent wash. Such remedies, applied to the eye by means of an eye-cup, or by wet pledgets, prove, doubtless, highly serviceable. Any of the undermentioned may, therefore, be used with this view; and where the pain is very acute, forty or fifty drops of the vinous tincture of opium may be added to any of them; or the eye may be frequently bathed with a decoction of bruised poppy-heads. In such cases a quarter of a grain of solid opium may also be taken internally, and repeated every four or six hours.*

For the purpose of allaying heat and inflammation of the eyes, some practitioners give the preference to warm instead of cold collyria; and among this number is Mr. Ware. The fact is, that inflammations of the eyes are known to yield sometimes to cold, and sometimes to warm fomentations. In cases of high irritation the warm may be used; but the alternate use of cold and hot applications has sometimes succeeded when neither of them singly appeared capable of putting an end to the diseased action.

When the disease has been of long standing, I have seen very extraordinary effects from a seton in the neck, or between the shoulders, especially the latter. It should be put upwards and downwards, or in the direction of the spine, and in the middle between the shoulder blades. It may be dressed twice a-day with yellow basilicon. I have known patients, who had been blind for a considerable time, recover sight by means of a seton placed as

* Take Sulphate of Zinc,	Rose Water,	6 ounces.
Supracetate of Lead, of each,	Mix them.	
8 grains.	or	
Mix them for an eye-wash.	Take Rose Water,	3 ounces.
or	Solution of the Acetate of Lead,	18 drops.
Take Solution of the Acetate of Ammonia,	Mix them.	
Rose Water, of each,	or	
2 ounces.	Take Sulphate of Zinc,	10 grains.
Camphor Mixture,	Rose Water,	
1 ounce.	Camphor Mixture, each,	3 ounces.
Mix them.	Mix them.	
or		
Take Purified Alum,		
½ drachm.		

above. When the seton is put across the neck, it soon wears out, and is both more painful and troublesome than between the shoulders; besides, it leaves a disagreeable mark, and does not discharge so freely.

When the heat and pain of the eyes are very great, a poultice of bread and milk, softened with sweet oil or fresh butter, may be applied to them, at least all night; and they may be bathed with lukewarm milk and water in the morning.

If the patient cannot sleep, which is sometimes the case, he may take twenty or thirty drops of laudanum, or two spoonsful of syrup of poppies, over-night, more or less according to his age, or the violence of the symptoms.

After the inflammation is gone off, if the eyes still remain weak and tender, they may be bathed every night and morning with cold water and a little brandy, six parts of the former to one of the latter. A method should be contrived by which the eye can be quite immersed in the brandy and water, where it should be kept for some time. I have generally found this, or cold water and vinegar, as good a strengthener of the eyes as any of the most celebrated collyria.*

When an inflammation of the eyes proceeds from a scrophulous habit, it generally proves very obstinate. In this case the patient's diet must not be too low, and he may be allowed to drink small negus, or now and then a glass of wine. The most proper medicine is the Peruvian bark, which may either be given in substance, or otherwise.

If ophthalmia be dependent on a venereal taint, mercury is the remedy to be depended upon for its removal. When it arises in a scrophulous habit, affecting the tarsi, or edges of the eye-lids, and is attended with ulcerations, as is often the case, Peruvian bark, with alteratives, mineral waters, and sea-bathing, will be the most proper remedies. In these cases the bark combined with hemlock has sometimes proved serviceable. At the same time that the use of these remedies is continued, the edges of the eyelids may be smeared, morning and night, with a little ointment,† composed of mercury, or the sulphate of zinc.

It will be proper frequently to look into the eyes, to see if any hairs be turned inwards, or pressing upon them.‡ These ought to be removed by plucking them out with a pair of small pincers.

* After the active stage of the inflammation has terminated, and the bloodvessels of the eye appear turgid and relaxed, excellent effects are often found to result from letting a drop or two of the tincture of opium fall into the eye. In this state the eye will bear the application of active stimuli with more advantage than is commonly believed.—In scrophulous inflammation of the eyes, sea-bathing, together with keeping the body open by gentle purgatives of sea-water, are eminently useful. I lately witnessed a case where the aqueous humour of the eye had become so turbid, that the patient, evidently of a scrophulous habit, could hardly distinguish light from darkness, which was almost wholly removed, and sight in great measure restored, by persisting in a course of sea-bathing during the months of summer. A. P. B.

† Take Ointment of the Nitrate of Mercury,

Prepared Lard, of each, $\frac{1}{2}$ ounce.

Make an ointment.

or
Take Sulphate of Zinc,

Prepared Lard,

Make an ointment.

1 scruple.

1 ounce.

‡ Any foreign body lodged in the eye may be expeditiously removed by passing a small hair pencil between the eyelid and the ball of the eye. In some places the peasants do this very effectually, by using their tongue in the same manner.

Those who are liable to frequent returns of this disease ought constantly to have an issue in one or both arms. Bleeding or purging in the spring and autumn will be very beneficial to such persons. They ought likewise to live with the greatest regularity avoiding strong liquor, and every thing of a heating quality. Above all, let them avoid the night air and late studies.* It may also be necessary, to prevent a return of ophthalmia, to continue the use of blisters behind the ears, or an issue or seton. The cold bath, employed by immersing the whole body, or by washing the head in cold water once or twice a-day, is also a powerful means of prevention. Also the application of cold water to the eyes themselves, or of any astringent collyrium, by means of an eye-cup, two or three times a-day, may likewise be serviceable in preventing its return, or removing it after it has become habitual. Tonics have also been adopted with this intention, and with the best effects.

CHAP. XX.

QUINSEY, OR INFLAMMATORY SORE THROAT. (*Cynanche Tonsillaris.*)

THIS disease is very common in Britain, and is frequently attended with great danger, occupying the glands, such as the tonsils, but frequently extending throughout the whole mucous membrane of the fauces, so as essentially to interrupt the speech, respiration, and deglutition of the patient. It prevails in the winter and spring, and is most fatal to young people of a sanguine temperament.

CAUSES.—In general it proceeds from the same causes as other inflammatory disorders, viz. exposure to cold, an obstructed perspiration, or whatever heats or inflames the blood. An inflammation of the throat is often occasioned by omitting some part of the covering usually worn about the neck, by drinking cold liquor when the body is warm, by riding or walking against a cold northerly wind, or any thing that greatly cools the throat and parts adjacent. It may likewise proceed from the neglect of bleeding, purging, or stoppage of any customary evacuation.

Singing, speaking loud and long, or whatever strains the throat, may likewise cause an inflammation of that organ. I have often known the quinsy prove fatal to jovial companions, who, after sitting long in a warm room, drinking hot liquors, and singing with vehemence, were so imprudent as to go abroad in the cold night air. Sitting with wet feet, or keeping on wet clothes, are very apt to occasion this malady. It is likewise frequently occasioned by continuing long in a moist place, sitting near an open window, sleeping in a damp bed, sitting in a room that has been

* As most people are fond of using eye-waters and ointments in this and other diseases of the eyes, we have inserted some of the most approved form of these medicines in the Appendix. See Appendix, *Eye-water* and *Eye-salve*.