

APPENDIX:

CONTAINING

1. A List of Simples and of such Medicinal Preparations as are commonly used in Practice, with their proper Doses.
2. The Method of preparing and compounding such Medicines as are recommended in the former Part of the Book, with the Addition of several others of a similar Nature.
3. Remarks on the Doses, Uses, and Manner of applying the different Preparations.

THE design of the following pages is, to exhibit such a list of drugs and medicines as may be necessary for private practice. They are considerably more numerous indeed than those recommended in the former part of the book, but are still greatly within the number contained in the most reformed dispensatories. The same medicine is seldom exhibited under different forms; and where different medicines answer nearly the same intention, there is commonly no more than one of them retained. Multiplying forms of medicine for the same intention tends rather to bewilder than assist the young practitioner, and the experienced physician can never be at a loss to vary his prescriptions as occasion requires. The chemical and other difficult preparations are for the most part omitted. All of them that are used by any private practitioner are not worth preparing. He will buy them much cheaper than he can make them. Great care, however, is necessary to obtain them genuine. They are often adulterated, and ought never to be purchased unless from persons of known veracity. Such of them as are in common use are inserted in the lists of drugs and medicines. Their proper doses and manner of application are mentioned in the practical part of the book, wherever they are prescribed.

Such articles of medicines as are to be found in the house or garden of almost every peasant, as barley, eggs, onions, &c. are likewise for the most part omitted. It is needless to swell a list of medicines with such things as can be obtained whenever they are wanted, and which spoil by being kept.

The preparations made and sold by distillers and confectioners are also generally left out. These people, by operating upon a larger plan, generally make things better, while it is in their power to afford them much cheaper than they can be prepared by any private hand.—The quantity ordered of every medicine is as small as could well be prepared, both to prevent unnecessary expense, and that the medicine might not spoil by keeping. Almost every medicine suffers by being kept, and should be used as soon after it has been prepared as possible. Even simple drugs are apt to spoil, and should therefore be laid in in small quantities; they either rot, are consumed by insects, or evaporate so as to lose their peculiar taste or flavour, and often become quite insignificant.

In several compositions, the ingredient on which the efficacy of the medicine principally depends is increased, while the auxiliaries, which are generally ordered in such trifling quantities as to be of no importance, are left out, or only such of them retained as are necessary to give the medicine a proper consistence, or the like.

The colouring ingredients are likewise for the most part omitted. They increase the bulk and price of the medicine; without adding any thing to its value. It would be well if they were never used at all. Medicines are often adulterated for the sake of a colour. Acrid and even poisonous substances are, for this purpose, sometimes introduced into those medicines which ought to be most bland and emollient. Ointment of elder, for example, is often mixed with verdigrise to give it a fine green colour, which entirely frustrates the intention of that mild ointment. Those who wish to obtain genuine medicines should pay no regard to their colour.

Some regard is likewise paid to expense. Such ingredients as greatly increase the price of any composition, without adding considerably to its virtue, are generally either omitted, or somewhat less expensive substituted in their place. Medicines are by no means powerful in proportion to their price. The cheapest are often the best; besides, they are the least apt to be adulterated, and are always most readily obtained.

With regard to the method of compounding medicines, I have generally followed that which seemed to be the most simple and natural, mentioning the different steps

of the process in the same order in which they ought to be taken, without paying an implicit regard to the method of other dispensaries.

I have followed the alphabetical order, both with regard to the simples and preparations. A more scientific method would have been agreeable to some persons, but less useful to the generality of readers. The different classes of medicine have no great dependence upon one another, and, where they have, it is hard to say which should stand first or last; no doubt the simple preparations ought to precede the more compound. But all the advantages arising from this method of arrangement do not appear equal to that single one, of being able, on the first opening of the book, to find out any article, which, by the alphabetical order, is rendered quite easy.

The dose of every medicine is mentioned, whenever it appeared necessary. When this is omitted, it is to be understood that the medicine may be used at discretion. The dose mentioned is always for an adult, unless when the contrary is expressed. It is not an easy matter to proportion the doses of medicine exactly to the different ages, constitutions, &c. of patients; but, happily for mankind, mathematical exactness here is by no means necessary.

Several attempts have been made to ascertain the proportional doses for the different ages and constitutions of patients; but, after all that can be said upon this subject, a great deal must be left to the judgment and skill of the person who administers the medicine. The following general proportions may be observed; but they are by no means intended for exact rules. A patient between twenty and fourteen may take two-thirds of the dose ordered for an adult; from fourteen to nine, one-half; from nine to six, one-third; from six to four, one-fourth; from four to two, one-sixth; from two to one, a tenth; and below one, a twelfth.

To prevent mistakes, the English name of every medicine is not only used, but the different articles are arranged according to the order of the English alphabet, and the smallest and largest dose placed opposite to the operation of each article. The doses indeed refer to adults, but may be adapted to different ages by attending to the rules laid down in the opposite page. Short cautions are occasionally inserted under such articles as require to be used with care.

Though a greater variety of medicines is contained in this than in any former edition of the Domestic medicine, yet the Author would advise those who peruse it, as far as possible, to adhere to simplicity in practice. Diseases are not cured by multiplicity of medicines, but by their proper application. A few simples, judiciously administered, and accompanied with a proper regimen, will do more good than a farrago of medicines employed at random.

OF THE

TICE,

WITH THEIR PARTICULAR VIRTUES AND PROPER DOSES.

*. * Explanation of the Abbreviations used in the following Doses: scr. scruple, scrs. scruples, dr. drachm, drs. drachms, oz. ounce, drps. drops, gr. grain, grs. grains, grad. gradually.

Names.	Properties.	Doses.
A CACIA, the expressed juice, from } Acid, the acetous - - - - - - , muriatic - - - - - - , nitrous, diluted - - - - - - , vitriolic, diluted - - - Ether, vitriolic - - - Ethiopi's mineral - - - Agaric, <i>used externally as a styptic, to staunch blood.</i> Aloes - - - Alum - - - - - - , burned - - - Amber, prepared - - - Ammoniac, gum - - - - - - , milk of - - - Angelica, the root powdered - - - Anise, the seeds - - - Antimony, crude - - - - - - , calcined - - - - - - , cinnamon of - - - - - - , glass of - - - - - - , tartarized - - -	Demulcent. Refrigerant, &c. Antiseptic, &c. Tonic, Febrifuge, &c. Tonic, Antiseptic. Anodyne. Alterative. Emenagogue, &c. Astringent. Escharotic, &c. Antispasmodic. Expectorant. Expectorant. Stimulant, &c. Carminative. Febrifuge, &c. Febrifuge. Diaphoretic. Emetic, never used. Emetic.	1 scr. to 1 dr. 1 scr. - 1 dr. 10 drps. - 40 drps. 15 drps. - 40 drps. 15 drps. - 40 drps. 30 drps. - 2 drs. 10 grs. - 30 grs. 5 grs. - 30 grs. 6 grs. - 20 grs. 3 grs. - 12 grs. 1 dr. - 1 dr. 6 grs. - 30 grs. 1 oz. - 1 oz. 1 dr. - 1 1/2 dr. 10 grs. - 1 dr. 10 grs. - 1 dr. 1 scr. - 1 dr. 10 grs. - 1 scr. 1 dr. - 2 grs. 1 gr. - 4 grs.
Asafoetida, - - - - - - , milk of - - - Asarum, - - -	{ Emenagogue, Expectorant, and Antispasmodic. Antispasmodic, &c. Emetic and Errhine.	grs. - 1 dr. 1 oz. - 1 oz. 3 grs. - 5 grs.
Balsam of capivi - - - - - - Canadian - - - - - - of Peru - - - - - - of Tolu - - - Bark, cascarilla - - - - - - , Peruvian, powder of - - - Bear's foot, powder of the leaves - - - Benzoin, resin of, not employed internally, and principally for obtaining Benzoic acid Bistort, powder of the root - - - Blessed Thistle - - - - - - , expressed, } - - - , juice of - - - Bole, Armenian - - - - - - , French - - - Borax, rarely used internally - - - Broom, ashes of the tops - - - Burdock, powder of the root - - -	Diuretic, &c. Diuretic, &c. Expectorant, Stimulant. Stimulant, Expectorant. Tonic, &c. Tonic. Narcotic. Astringent. Tonic, Emetic. <i>The same.</i> Astringent. <i>The same.</i> Detergent. Diuretic. Sudorific, &c.	20 drps. - 60 drps. 1 scr. - 1 dr. - - - 15 grs. - 2 scrs. 10 grs. - 1 dr. 1 scr. - 2 drs. 10 grs. - 20 grs. 1 scr. - 1 dr. 10 grs. - 1 dr. 2 drs. - 2 oz. 10 grs. - 2 drs. - - - 10 grs. - 40 grs. 1 scr. - 1 dr. 10 grs. - 1 dr.
Calamine stone, levigated, <i>used externally in Turner's cerate.</i> Calomel - - -	{ Alterative. Purgative.	1 gr. - 3 grs. 3 grs. - 12 grs.

Names.	Properties.	Doses.
Camphor	Narcotic, Diaphoretic.	2 grs. to $\frac{1}{2}$ scr.
Canella alba, powder of	Tonic, Carminative.	1 scr. to 2 drs.
Cantharides	Stimulant, Diuretic.	$\frac{1}{2}$ gr. - 4 grs.
Cardamoms	Carminative.	5 grs. - 20 grs.
Caraway seeds	Carminative.	10 grs. - 40 grs.
Carrot, seed of the wild	Carminative.	1 scr. - 1 dr.
Cascarilla bark	Tonic.	10 grs. - 40 grs.
Cassia, the pulp	Laxative.	2 drs. - 1 oz.
Castor	Antispasmodic.	8 grs. - 1 dr.
Caustic, lunar; used <i>externally</i> as an escharotic; <i>internally</i> , gr. $\frac{1}{2}$, cautiously increased to grs. ij, in epilepsy.		
Catechu	Astringent.	15 grs. - 30 grs.
Camomile, in powder	Tonic.	20 grs. - 1 dr.
Centaur, the lesser	Tonic.	1 scr. - 1 dr.
Chalk	Absorbent.	20 grs. - 2 scrs.
Cinnamon	Carminative.	5 grs. - 1 dr.
Colocynth	Violently Cathartic.	2 grs. - 10 grs.
Columbo	Tonic.	10 grs. - 1 dr.
Confection, aromatic	Cordial.	10 grs. - 2 scrs.
" opiate	Anodyne.	10 grs. - 2 scrs.
Crab's claws, prepared	Absorbent.	10 grs. - 1 dr.
Conserve of roses	Astringent.	1 dr. - 1 oz.
Contrainerva	Febrifuge.	10 grs. - 2 scrs.
Coriander seed	Carminative.	15 grs. - 1 dr.
Cowhage,	Vermifuge.	{ The spiculæ of one pod mixed with honey or molasses.
Cummin seed	Stimulant.	
Dandelion, expressed juice of	Diuretic.	1 oz. - 3 oz.
Decoction of hartshorn	Demulcent.	{ Half a pint, repeated as often as necessary.
" of broom tops	Diuretic.	
" of Peruvian bark	Tonic.	{ 1 oz. to a pint of water; to be taken by teacupsful.
" of the inner bark of the elm	Diuretic.	
" of sarsaparilla	Alterative & diaphoretic.	4 oz. - 10 oz. daily.
Decoction of sarsaparilla, compound	Alterative & diaphoretic.	4 oz. - 16 oz. daily.
" of guaiacum	Diaphoretic.	{ 3 drachms to a pint of water. A pint daily.
Deadly night shade	Narcotic, gr. $\frac{1}{2}$ to grs. ij.	
Dragon's blood	Astringent.	10 grs. - 2 scrs.
Earth, fuller's, use external in excoriations	Astringent.	
Electuary of cassia	Aperient.	1 dr. - 1 oz.
" of scammony	Purgative.	20 grs. - 2 drs.
" lenitive, or of senna	Purgative.	30 grs. - 6 drs.
Elixir of vitriol	Tonic.	15 drps. - 50 drps.
Elecampane, powder of the root	Stimulant.	20 grs. - 1 dr.
Extract of broom-tops	Diuretic.	$\frac{1}{2}$ dr. - 1 dr.
" Peruvian bark	Tonic.	10 grs. - $\frac{1}{2}$ dr.
" cascarilla	Tonic.	10 grs. - $\frac{1}{2}$ dr.
" camomile	Tonic.	10 grs. - 1 dr.
" colocynth comp.	Cathartic.	5 grs. - 25 grs.
Extract of gentian	Stomachic.	10 grs. - $\frac{1}{2}$ dr.
" hemlock	Alterative.	2 grs. - 10 grs.
" liquorice	Demulcent.	1 dr. - $\frac{1}{2}$ oz.
" logwood	Astringent.	10 grs. - $\frac{1}{2}$ dr.
" black hellebore	Emenagogue.	3 grs. - 10 grs.
" jalap	Purgative.	5 grs. - 20 grs.
" guaiacum	Diaphoretic.	10 grs. - 20 grs.
" white poppies	Anodyne.	1 gr. - 5 grs.
" rue	Emenagogue.	10 grs. - 20 grs.
" savin	The same.	10 grs. - 30 grs.
" senna	Aperient.	10 grs. - 30 grs.
" wormwood	Tonic.	10 grs. - 30 grs.
Fern, powder of the root	Vermifuge.	$\frac{1}{2}$ dr. - $\frac{1}{2}$ oz.

Names.	Properties.	Doses.
Fennel seed	Aromatic.	20 grs. to 1 dr.
Foxglove, powder of the leaves	Diuretic.	$\frac{1}{2}$ gr. - 3 grs.
	or a drachm infused in a pint of boiling water, of which a dose is	1 oz.
	(Administered with caution.)	
Frankincense		10 grs. - 30 grs.
Flowers of camomile, powder of	Tonic.	10 grs. - 1 dr.
elder	Cathartic, &c.	
rosemary	Emenagogue.	
damask roses, rarely employed unless to make rose-water.	Laxative.	
red ditto, in infusion	Astringent.	Ad libitum.
Fruits. Almonds	Demulcent.	
Figs, dried	Aperient.	
French prunes	Aperient.	
Tamarinds	Aperient.	
Galbanum	Deobstruent.	10 grs. - $\frac{1}{2}$ dr.
Galls	Astringent.	10 grs. - 20 grs.
Garlic, cloves of	Expectorant.	No. 1. - No. 6.
Gentian	Tonic.	10 grs. - 40 grs.
Germander	Tonic.	15 grs. - 1 dr.
Ginger	Carminative.	5 grs. - 20 grs.
Ginseng		20 grs. - 30 grs.
Guaiacum, wood of	Diaphoretic.	
gum-resin	The same.	10 grs. - 30 grs.
Gum-arabic	Demulcent.	15 grs. - 1 dr.
gamboge	Hydragogue.	2 grs. - 12 grs.
Hartshorn, prepared	Emollient.	20 grs. - 1 dr.
spirits of		10 drps. - 40 drps.
Hellebore, black	Emenagogue.	5 grs. - 10 grs.
white	Emetic, &c.	1 gr. - 5 grs.
Hemlock	Narcotic.	Should always be begun with very small doses, as one grain or less, and gradually increased as the constitution will bear. See extract of.
Hiera picra	Purgative.	10 grs. - 20 grs.
Honey of squills	Diuretic.	10 grs. - 40 grs.
of roses	Astringent.	1 dr. - 2 drs.
Hoffman's anodyne liquor	Anodyne, &c.	20 drps. - 60 drps.
Infusion of gentian, compound	Tonic.	1 oz. - 3 oz.
roses	Astringent.	2 oz. - 8 oz.
senna	Aperient.	$\frac{1}{2}$ oz. - 2 oz.
Ipecacuanha	Emetic, and expectorant, gr. i. to iij.	10 grs. - 30 grs.
Iris, florentine	Tonic.	1 scr. - 1 dr.
Iron, rust of	Tonic.	5 grs. - 20 grs.
ammoniated	The same.	2 grs. - 10 grs.
tartarized	Tonic.	2 grs. - 10 grs.
Jalap, powder of	Purgative.	10 grs. - 40 grs.
Juniper, powder of the berries	Diuretic.	20 grs. - 1 dr.
Kino, gum	Astringent.	10 grs. - 30 grs.
Lead, white	Astringent.	$\frac{1}{2}$ gr. - 2 grs.
sugar of	Demulcent.	10 grs. - 40 grs.
Lichen, ash-coloured, ground	The same.	1 oz. - 4 oz.
Icelandic, a strong decoction of	Refrigerant.	4 oz. - 8 oz.
Lime-water	Lethontripic.	15 drps. - 40 drps.
Lixivium of tartar		An infusion of 1 ounce to a quart of water, may be used at pleasure.
Linseed	Demulcent.	$\frac{1}{2}$ dr. - 1 dr.
Liquorice, root of	Demulcent.	

<i>Names.</i>	<i>Properties.</i>	<i>Doses.</i>
Madder powder - - -	- - -	$\frac{1}{2}$ dr - 1 dr.
Mace - - -	Stomachic.	10 grs. - 20 grs.
Magnesia - - -	Antacid.	$\frac{1}{2}$ dr. - 2 drs.
——, calcined - - -	<i>The same.</i>	—
Manna - - -	Aperient.	$\frac{1}{2}$ oz. - 2 oz.
Marsh-mallows, root and leaves } of }	Demulcent.	$\frac{1}{2}$ dr. - 1 dr.
Mastich, gum - - -	Carminative.	10 grs. - 30 grs.
Mercury, crude - - -	- - -	- - -
——, calcined - - -	Alternative.	$\frac{1}{2}$ gr. - 2 grs.
——, with chalk - - -	Alternative, Antisyphilitic.	10 grs. - 30 grs.
——, corrosive sublimate - - -	<i>The same.</i>	$\frac{1}{2}$ gr. - $\frac{1}{2}$ gr.
——, cinnabar of - - -	Alternative, &c.	10 grs. - 30 grs.
——, red precipitate of } ——, white ditto }	Use chiefly external.	- - -
——, yellow emetic - - -	Sternutatory.	1 gr. - 3 grs.
Mezereon - - -	Sialogogue in Decoct.	To a pint of water, 2 drs.
Millipedes - - -	Expectorant.	20 grs. - 2 drs.
Musk - - -	Antispasmodic.	5 grs. - 40 grs.
Mustard seed - - -	Stimulant. Emetic.	1 dr. - 1 oz.
Myrrh, gum - - -	Expectorant.	10 grs. - 1 dr.
Nitre, purified - - -	Diuretic.	10 grs. - 30 grs.
Nutmeg - - -	Stomachic.	6 grs. - $\frac{1}{2}$ dr.
Oil of Almonds - - -	Demulcent.	$\frac{1}{2}$ oz. - 1 oz.
—— amber, rectified - - -	Antispasmodic.	10 drps. - 30 drps.
—— aniseed - - -	Carminative.	1 drp. - 5 drps.
—— castor - - -	- - -	2 drs. - 1 oz.
—— cinnamon - - -	Stimulant.	1 drp. - 3 drps.
—— juniper - - -	Diuretic.	2 drps. - 10 drps.
—— lemon peel - - -	- - -	2 drps. - 5 drps.
—— linseed - - -	Demulcent.	$\frac{1}{2}$ oz. - 1 oz.
—— olive - - -	Demulcent, Laxative.	$\frac{1}{2}$ oz. - 1 oz.
—— palm - - -	Use external.	- - -
—— peppermint - - -	Stimulant, &c.	1 drp. - 3 drps.
—— turpenne - - -	{ Diuretic. Externally, } stimulant. }	10 drps. - 30 drps.
Onion, expressed juice of - - -	A powerful diuretic.	$\frac{1}{2}$ oz. - 2 oz.
Opium - - -	Narcotic.	$\frac{1}{4}$ gr. - 2 grs.
Opoponax - - -	Emenagogue.	10 grs. - 30 grs.
Oyster-shells, prepared - - -	Absorbent.	$\frac{1}{2}$ dr. - 2 drs.
Oxymel of colchicum - - -	Expectorant.	$\frac{1}{2}$ dr. - 1 dr.
—— of squills - - -	Diuretic.	$\frac{1}{2}$ dr. - 1 dr.
Pennyroyal - - -	Emenagogue.	- - -
Peppermint - - -	Stimulant.	- - -
Petroleum - - -	Antispasmodic.	10 drps. - 30 drps.
Pills, aloetic - - -	Purgative.	10 grs. - 30 grs.
——, of the gums - - -	Diuretic.	10 grs. - 30 grs.
——, mercurial - - -	Antisyphilitic.	10 grs. - 20 grs.
Pitch, Burgundy - - -	- - -	- - -
Pomegranate, powder of - - -	- - -	20 grs. - 1 dr.
Poppy heads - - -	Anodyne.	- - -
Powder, antimonial - - -	Febrifuge.	3 grs. - 6 grs.
May be taken according to the directions for James's powder, with which, in effects it nearly coincides.		
—— of contrayerva, compound - - -	<i>The same.</i>	15 grs. - 30 grs.
—— of chalk, compound - - -	Absorbent.	20 grs. - 40 grs.
——, with opium - - -	Absorbent and anodyne.	10 grs. - 40 grs.
Powder of ipecacuanha, com- } pound, or Dover's powder. }	Diaphoretic.	10 grs. - 30 grs.
Quassia - - -	Tonic.	5 grs. - 30 grs.
(One drachm to a pint of water for an infusion.)		
Quince seeds, mucilage of, - - -	Demulcent. At pleasure, to obtund acrimony.	- - -
Rhubarb, powder of - - -	Purgative.	10 grs. - 40 grs.
Resin, yellow - - -	Diuretic.	3 grs. - 20 grs.
Rue powder - - -	Emenagogue.	20 grs. - 40 grs.
Sassafras - - -	Diaphoretic, Altera- } tive, &c. }	1 scr. - 1 dr.

Names.	Properties.	Doses.
Savin	Emenagogue.	10 grs. - 30 grs.
St. John's wort	Diaphoretic.	20 grs. - 1 dr.
Saffron	Cordial, &c.	5 grs. - 20 grs.
Sagapenum		10 grs. - 30 grs.
Sal ammoniac	Stimulant.	10 grs. - 30 grs.
Salt, Epsom	Aperient.	2 drs. - 1 oz.
—, Glauber	Aperient.	4 drs. - 2 oz.
—, of hartshorn	Cordial.	10 grs. - 20 grs.
—, Polychrest	Aperient.	20 grs. - 1 oz.
—, Rochel	Aperient.	2 scrs. - 1 oz.
—, of tartar	Aperient.	10 grs. - 30 grs.
Saunders, red	Astringent.	1 dr. - 1 dr.
Sarsaparilla, powder of	Alterative.	20 grs. - 40 grs.
Scammony	Cathartic.	5 grs. - 20 grs.
Seneka	Diaphoretic.	20 grs. - 40 grs.
Senna	Aperient.	20 grs. - 40 grs.
Soap	Lithontriptic.	20 grs. - 1 oz.
—, lees	The same.	10 drps. - 30 drps.
Spearmint	Stimulant. Cordial.	10 grs. - 2 scrs.
Senna	Aperient.	1 scr. - 1 dr.
Scurvy-grass, expressed juice	Antiscorbutic.	1 oz. - 4 oz.
Snake-root	Diaphoretic.	20 grs. - 40 grs.
Sorrel, juice of, depurated	Antiscorbutic.	2 oz. - 4 oz.
Spirit of lavender	Cordial, Stimulant.	1 dr. - 2 drs.
—, of mindererus	Diaphoretic.	1 dr. - 1 oz.
—, sweet, of vitriol	Tonic.	15 drps. - 40 drps.
—, of nitre	Diuretic.	15 drps. - 40 drps.
—, of sal ammoniac	Diaphoretic, &c.	15 drps. - 40 drps.
—, compound		—
—, fetid	Antispasmodic, &c.	—
Steel, filings of	Tonic, Emenagogue.	5 grs. - 1 scr.
Spermaceti	Demulcent.	20 grs. - 1 dr.
Sponge, burned	Deobstruent.	20 grs. - 1 dr.
Storax	Astringent, &c.	10 grs. - 1 dr.
Sulphur	Cathartic and diaphoretic.	20 grs. - 1 dr.
—, precipitated, of antimony	Alterative.	1 gr. - 4 grs.
Squill, dried powder	Diuretic.	1 gr. - 3 grs.
—, fresh	The same.	5 grs. - 15 grs.
Syrup of poppies	Anodyne.	1 dr. - 1 oz.
—, of buckthorn	Cathartic.	1 dr. - 2 drs.
—, of ginger	Carminative.	1 dr. - 1 oz.
—, of meadow saffron	Diuretic, &c.	1 dr. grad. to 1 oz.
—, of violets	Gently laxative.	1 dr. - 2 drs.
—, of poppies	Anodyne. To children,	1 dr. - 2 oz.
—, s in general		1 dr. - 2 drs.
Tar	Pulmonary.	5 grs. - 1 scr.
—, water of		A pint daily.
Tartar, cream of	Refrigerant and aperient.	2 drs. - 1 oz.
—, emetic	Alterative.	1 gr. - 1 gr.
	An Emetic.	1 gr. - 3 grs.
Terra japonica	Astringent.	20 grs. - 40 grs.
Tobacco, an infusion of, 1 drachm	to a pint of water; should be administered by table	
spoonsful; powerfully diuretic;	or in form of clyster.	
Tin, powder of	Anthelmintic.	20 grs. - 1 dr.
Turmeric		20 grs. - 1 dr.
Turpentine, spirit of	Diuretic.	10 drps. - 30 drps.
—, Venice	The same.	20 grs. - 1 dr.
Tincture of aloes	Purgative.	1 oz. - 1 oz.
—, compound	The same.	1 dr. - 2 drs.
—, of asafetida	Antispasmodic.	1 dr. - 2 drs.
—, of Benzoin, compound	Vulnerary.	10 drps. - 40 drps.
—, of cantharides	Diuretic.	10 drps. - 40 drps.
—, of cardamoms	Carminative	1 dr. - 1 oz.
—, of castor		1 dr. - 1 dr.
—, of catechu	Astringent.	1 dr. - 2 drs.
—, of Peruvian bark	Tonic.	1 dr. - 1 oz.
—, of iron, muriated	Tonic.	10 drps. - 60 drps.
—, of Calumbo	Tonic.	1 dr. - 3 drs.
—, of gentian, compound	Tonic.	1 dr. - 3 drs.
—, of guaiacum volatile	Diaphoretic.	1 dr. - 3 drs.
—, of black hellebore	Emenagogue.	1 scr. - 1 dr.

<i>Names.</i>	<i>Properties.</i>	<i>Doses.</i>
Tincture of jalap - - -	Purgative.	1 dr. to ½ oz.
— of myrrh - - -	Emenagogue.	1 scr. - 1 dr.
— of opium - - -	Narcotic.	10 drps. - 40 drps.
— camphorated, or paregoric elixir - - -	Anodyne and antispasmodic.	1 dr. - 3 drs.
— of rhubarb - - -	Purgative, Stomachic.	½ oz. - 2 oz.
— of senna - - -	Cathartic, Stomachic.	2 drs. - 1 oz.
— of snake-root - - -	Diaphoretic, &c.	1 dr. - 2 drs.
— of valerian - - -	Antispasmodic.	1 dr. - 3 drs.
— volatile - - -	The same and stimulant.	½ dr. - 2 drs.
Tormentil, powder of - - -	Astringent.	10 grs. - 1 dr.
Tumeric - - -	—	1 scr. - 1 dr.
Turpentine, Venice - - -	Diuretic and cathartic.	1 scr. - 1 dr.
Tutty, levigated - - -	Used externally in ointments and cerates.	
Uva ursi, in powder - - -	Lithontriptic.	20 grs. - 1 dr.
Valerian, wild, powder of - - -	Stimulant, Antispasmodic.	20 grs. - 2 drs.
Vinegar, distilled - - -	Refrigerant, Antiseptic.	2 drs. - 1 oz.
— of squills - - -	Diuretic.	10 drps. - 50 drps.
—	As an emetic.	½ oz. - 1 oz.
Verdigris - - -	Violently emetic.	1 gr. - 2 grs.
Vitriol, white - - -	As a Tonic.	2 grs. - 5 grs.
—	As a quickly operating emetic.	20 grs. - 1 dr.
—, blue - - -	Emetic.	1 gr. - 3 grs.
Watercress, expressed juice of - - -	Antiscorbutic.	½ oz. - 2 oz.
— trefoil - - -	—	½ dr. - 1 dr.
Waters, any of the simple distilled - - -	Used as vehicles.	½ oz. - 4 oz.
Wax, white - - -	Demulcent and emollient.	20 grs. - 1 dr.
—, yellow - - -	—	½ oz. - 2 oz.
Wormwood, expressed juice - - -	Vermifuge, Tonic.	1 gr. - 5 grs.
White lead - - -	Astringent.	½ gr. to grs. ij. of the extract.
Wild cucumber - - -	Cathartic, Hydragogue.	½ oz. - 1 oz.
Wine, aloetic - - -	Purgative.	20 drps. - 2 drs.
—, antimonial - - -	Emetic.	1 dr. - ½ oz.
—, ipecacuanha - - -	Emetic.	½ oz. - 2 oz.
—, rhubarb - - -	Purgative.	½ oz. - 2 oz.
Zedoary - - -	Stomachic.	10 grs. - 40 grs.
Zinc, flowers of - - -	Tonic. Antispasmodic.	3 grs. - 10 grs.

DOMESTIC PHARMACOPŒIA.

MEDICINAL PREPARATIONS.

BALSAMS.

THE subject of this section is not the natural balsams, but certain compositions, which, from their being supposed to possess the balsamic qualities, generally go by that name.

This class of medicines was formerly very numerous, and held in great esteem. Modern practice, however, has justly reduced it to a very narrow compass.

Anodyne Balsam.—Take of white Spanish soap, one ounce; opium, unprepared, two drachms; rectified spirit of wine, nine ounces. Digest them together in a gentle heat for three days; then strain off the liquor, and add to it three drachms of camphor.

This balsam is intended to ease pain. It is of service in violent strains and rheumatic complaints, when not attended with inflammation. It must be rubbed with a warm hand on the part affected; or a linen rag moistened with it may be applied to the part, and renewed every third or fourth hour, till the pain abates. If the opium is left out, this will be the *Saponaceous Balsam*.

Locatelli's Balsam.—Take of olive oil, one pint; Strasburg turpentine and yellow wax, of each half a pound; red saunders, six drachms. Melt the wax with some part of the oil, over a gentle fire; then adding the remaining part of the oil and the turpentine; afterwards mix in the saunders, previously reduced to a powder, and keep them stirring together till the balsam is cold.

This balsam is recommended in erosions of the intestines, the dysentery, hæmorrhages, internal bruises, and in some complaints of the breast. Outwardly it is used for healing and cleansing wounds and ulcers. The dose when taken internally, is from two scruples to two drachms.

The vulnerary Balsam. (Tincture of Benjamin.)—Take of benzoin, powdered, three ounces; balsam of Peru, two ounces; hepatic aloes, in powder, half an ounce; rectified spirits of wine, two pints. Digest them in a gentle heat for three days, and then strain the balsam.

This balsam, or rather tincture, is applied externally to heal recent wounds and bruises. It is likewise employed internally to remove coughs, asthmas, and other complaints of the breast. It is said to ease the colic, cleanse the kidneys, and to heal internal ulcers, &c. The dose is from twenty to sixty drops.

This, though a medicine of some value, does not deserve the extravagant encomiums which have been bestowed on it. It has been celebrated under the different names of *The Commander's Balsam*, *Persian Balsam*, *Balsam of Berne*, *Wade's Balsam*, *Friar's Balsam*, *Jesuit's Drops*, *Turlington's Drops*, &c.

BOLUSES.

As boluses are intended for immediate use, volatile salts, and other ingredients improper for being kept, are admitted into their composition. They are generally composed of powders, with a proper quantity of syrup, conserve, or mucilage. The lighter powders are commonly made up with syrup, and the more ponderous, as mercury, &c., with conserve; but those of the lighter kind would be more conveniently made up with mucilage, as it increases their bulk less than the other additions, and likewise occasions the medicine to pass down more easily.

Astringent Bolus.—Take of alum, in powder, fifteen grains; gum kino, five grains; syrup, a sufficient quantity to make a bolus.

In an excessive flow of the menses, and other violent discharges of blood, proceeding from relaxation, this bolus may be given every four or five hours, till the discharge abates.

Diaphoretic Bolus.—Take of gum guaiacum, in powder, ten grains; flowers of sulphur and cream of tartar, of each one scruple; simple syrup, a sufficient quantity.

In rheumatic complaints, and disorders of the skin, this bolus may be taken twice a-day. It will also be of service in the inflammatory quinsy.

Mercurial Bolus.—Take of calomel, six grains; conserve of roses, half a drachm. Make a bolus.

Where mercury is necessary, this bolus may be taken twice or thrice a-week. It may be taken over-night; and if it does not operate, a few grains of jalap, or half an ounce of Epsom salts, will be proper next day to carry it off.

Bolus of Rhubarb and Mercury.—Take of the best rhubarb, in powder, from a scruple to half a drachm; of calomel, from four to six grains; simple syrup, a sufficient quantity to make a bolus.

This is a proper purge in hypochondriac constitutions; but its principal intention is to expel worms. Where a stronger purge is necessary, jalap may be used instead of the rhubarb.

Pectoral Bolus.—Take of spermaceti, a scruple; gum ammoniac, ten grains; salt of hartshorn, six grains; simple syrup, as much as will make them into a bolus.

This bolus is given in colds and coughs of long standing, asthmas, and beginning consumptions of the lungs. It is generally proper to bleed the patient before he begins to use it.

Purging Bolus.—Take of jalap, in powder, a scruple; cream of tartar, two scruples. Let them be rubbed together, and formed into a bolus, with simple syrup.

Where a mild purge is wanted, this will answer the purpose very well. If a stronger dose is necessary, the jalap may be increased to half a drachm or upwards.

CATAPLASMS AND SINAPISMS.

CATAPLASMS possess few or no virtues superior to a poultice, which may be so made, as, in most cases, to supply their place. They are chiefly intended either to act as discutients, or to promote suppuration; and as they may be of service in some cases, we shall give a specimen of each kind.

Discutient Cataplasm.—Take of barleymeal, six ounces; fresh hemlock leaves, bruised, two ounces; vinegar, a sufficient quantity. Boil the meal and hemlock in the vinegar for a little time, and then add two drachms of the sugar of lead.

Ripening Cataplasm.—Take of white lily-root, four ounces; fat figs and raw onions bruised, of each one ounce; yellow basilicum ointment, two ounces; gum galbanum, half an ounce; linseed-meal, as much as necessary. Boil the roots along with the figs in a sufficient quantity of water; then bruise and add to them the other ingredients, so as to form the whole into a soft cataplasm. The galbanum must be previously dissolved with the yolk of an egg.

Where it is necessary to promote suppuration, this cataplasm may be used by those who choose to be at the trouble and expense of making it. For my part, I have never found any application more proper for this purpose than a poultice of bread and milk, with a sufficient quantity of either boiled or raw onion in it, and softened with oil or fresh butter.

Sinapisms.—Sinapisms are employed to recal the blood and spirits to a part, as in the palsy and atrophy. They are also of service in deep-seated pains, as the sciatica, &c. When the gout seizes the head or the stomach, they are applied to the feet to bring the disorder to those parts. They are likewise applied to the patient's soles in the low state of fevers. They should not be suffered to lie on, however, till they have raised blisters, but till the parts become red, and will continue so when pressed with the finger.

The sinapism is only a poultice made with vinegar instead of milk, and rendered warm and stimulating by the addition of mustard, horse-radish, or garlic. The common sinapism is made by taking crumb of bread and mustard-seed in powder, of each equal quantities; strong vinegar, as much as is sufficient, and mixing them so as to make a poultice. When sinapisms of a more stimulating nature are wanted, a little bruised garlic may be added to the above.

CLYSTERS.

THIS class of medicines is of more importance than is generally imagined. Clysters serve not only to evacuate the contents of the belly, but also to convey very active medicines into the system. Opium, for example, may be administered in this way when it will not sit upon the stomach, and also in larger doses than at any time it can be taken by the mouth. The Peruvian bark may likewise be, with good effect, administered in form of clyster to persons who cannot take it by the mouth. A simple clyster can seldom do hurt, and there are many cases where it may do much good. A clyster even of warm water, by serving as a fomentation to the parts, may be of considerable service in inflammations of the bladder, and the lower intestines, &c.

Some substances, as the smoke of tobacco, may be thrown into the bowels in this

way, which cannot be done by any other means whatever. This may be easily effected by means of a pair of hand-bellows, with an apparatus fitted to them for that purpose.

The use of clysters is not merely confined to medicines. Aliment may also be conveyed in his way. Persons unable to swallow, have been, for a considerable time, supported by clysters.

Emollient Clyster.—Take of linseed-tea and new milk, each six ounces. Mix them. If fifty or sixty drops of laudanum be added to this, it will supply the place of the *Anodyne Clyster*.

Laxative Clyster.—Take of milk and water, each six ounces; sweet oil or fresh butter, and brown sugar, of each two ounces. Mix them.

If an ounce of Glauber's salt, or two table-spoonsful of common salt, be added to this, it will be the *Purging Clyster*.

Carminative Clyster.—Take of camomile flowers, an ounce; anise-seeds, half an ounce. Boil in a pint and a half of water to one pint.

In hysteric and hypochondriac complaints, this may be administered instead of the *Fetid Clyster*, the smell of which is so disagreeable to most patients.

Oily Clyster.—To four ounces of the infusion of camomile flowers, add an equal quantity of Florence oil.

This clyster is beneficial in bringing off the small worms lodged in the lower parts of the alimentary canal. When given to children, the quantity must be proportionably lessened.

Starch Clyster.—Take jelly of starch, four ounces; linseed oil, half an ounce. Liquefy the jelly over a gentle fire, and then mix in the oil.

In the dysentery or bloody flux, this clyster may be administered after every loose stool, to heal the ulcerated intestines, and blunt the sharpness of corroding humours. Forty or fifty drops of laudanum may be occasionally added; in which case, it will generally supply the place of the *Astringent Clyster*.

Turpentine Clyster.—Take of common decoction, ten ounces; Venice turpentine, dissolved with the yolk of an egg, half an ounce; Florence oil, one ounce. Mix them.

This diuretic clyster is proper in obstructions of the urinary passages, and in colicky complaints, proceeding from gravel.

Vinegar Clyster.—This clyster is made by mixing three ounces of vinegar with five of water-gruel. It answers all the purposes of a common clyster, with the peculiar advantage of being proper either in inflammatory or putrid disorders, especially in the latter.

COLLYRIA, OR EYE-WATERS.

EYE-WATERS have been multiplied without number, almost every person pretending to be possessed of some secret preparation for the cure of sore eyes. I have examined many of them, and find that they are pretty much alike, the basis of most of them being either alum, vitriol, or lead. Their effects evidently are to brace and restore the tone of the parts: hence they are principally of service in slight inflammations; and in that relaxed state of the parts which is induced by obstinate ones.

Camphor is commonly added to these compositions; but as it seldom incorporates properly with the water, it can be of little use. Boles and other earthy substances, as they do not dissolve in water, are likewise unfit for this purpose.

Collyrium of Alum.—Take of alum half a drachm; agitate it well together with the white of an egg.

This is the Collyrium of Riverius. It is used in inflammation of the eyes, to allay heat, and restrain the flux of humours. It must be spread upon linen, and applied to the eyes; but should not be kept on above three or four hours at a time.

Vitriolic Collyrium.—Take of white vitriol, half a drachm; rose-water, six ounces. Dissolve the vitriol in the water, and filter the liquor.

This, though simple, is perhaps equal in virtue to most of the celebrated collyria. It is an useful application in weak, watery, and inflamed eyes. Though the slighter inflammations will generally yield to it, yet in those of a more obstinate nature the assistance of bleeding and blistering will often be necessary.

When a strong astringent is judged proper, a double or triple quantity of the vitriol may be used. I have seen a solution of four times the strength of the above used with manifest advantage.

Collyrium of Lead.—Take sugar of lead, and crude sal ammoniac, of each four

grains. Dissolve them in eight ounces of common water : to which forty or fifty drops of laudanum may be occasionally added.

Those who choose, may substitute, instead of this, the collyrium of lead, recommended by Goulard ; which is made by putting twenty-five drops of his *extract of lead* to eight ounces of water, and adding a tea-spoonful of brandy. Indeed, common water and brandy, without any other addition, will in many cases answer very well as a collyrium. An ounce of the latter may be added to five or six ounces of the former ; and the eyes, if weak, bathed with it night and morning. For a rheum in the eyes, great benefit has been found by washing them frequently with rose-water. Many experience the like good effect from anointing the eye-lids with Smellon's salve.

CONFECTIONS.

CONFECTIONS, containing above sixty ingredients, are still to be found in some of the most reformed dispensaries. As most of their intentions, however, may be more certainly, and as effectually answered by a few glasses of wine or grains of opium, we shall pass over this class of medicines very slightly.

Japonic Confection, or Confection of Catechu.—Take of Japan earth, three ounces ; tormentil root, nutmeg, olibanum, of each two ounces ; opium dissolved in a sufficient quantity of Lisbon wine, a drachm and a half ; simple syrup and conserve of roses, of each fourteen ounces. Mix, and make them into an electuary. It supplies the place of the *Diascordium*. The dose is from a scruple to a drachm.

CONSERVES AND PRESERVES.

THESE preparations possess very few medical properties, and may rather be classed among sweetmeats than medicines. They are sometimes, however, of use, for reducing into boluses or pills some of the more ponderous powders, as the preparations of iron, mercury, and tin.

Conserves are compositions of fresh vegetables and sugar, beaten together into a uniform mass. In making these preparations, the leaves of vegetables must be freed from their stalks, the flowers from their cups, and the yellow part of orange-peel taken off with a rasp. They are then to be pounded in a marble mortar, with a wooden pestle, into a smooth mass ; after which, thrice their weight of fine sugar is commonly added by degrees, and the beating continued till they are uniformly mixed ; but the conserve will be better if only twice its weight of sugar be added.

Those who prepare large quantities of conserve generally reduce the vegetables to a pulp by the means of a mill, and afterwards beat them up with sugar.

Conserve of Red Roses.—Take a pound of red rose-buds, cleared of their heels ; beat them well in a mortar, and adding by degrees two pounds of double-refined sugar, in powder, make a conserve. After the same manner are prepared the conserves of orange-peel, rosemary-flowers, sea-wormwood, the leaves of wood-sorrel, &c.

The conserve of roses is one of the most agreeable and useful preparations belonging to this class. A drachm or two of it dissolved in warm milk, is ordered to be given as a gentle restringent in weakness of the stomach, and likewise phthisical coughs and spitting of blood. To have any considerable effects, however, it must be taken in large quantities.

Conserve of Sloes.—This may be made by boiling sloes gently in water, being careful to take them out before they burst ; afterwards expressing the juice, and beating it up with three times its weight of fine sugar.

In relaxations of the *uvula* and glands of the throat, this makes an excellent gargle, and may be used at discretion.

Preserves are made by steeping or boiling fresh vegetables first in water, and afterwards in syrup, or a solution of sugar. The subject is either preserved moist in the syrup, or taken out and dried, that the sugar may candy upon it. The last is the most useful method.

Candied Orange-Peel.—Soak Seville orange-peel in several waters, till it loses its bitterness ; then boil it in a solution of double-refined sugar in water, till it becomes tender and transparent.

Candied lemon-peel is prepared in the same manner.

DECOCTIONS.

Water readily extracts the gummy and saline parts of vegetables ; and though its action is chiefly confined to these, yet the resinous and oily being intimately blended with the gummy and saline, are in great part taken up along with them. Hence watery decoctions and infusions of vegetables constitute a large, and not unuseful, class of medicines. Although most vegetables yield their virtues to water, as well by infusion as decoction, yet the latter is often necessary, as it saves time, and does in a few minutes what the other would require hours, and sometimes days, to effect.

The medicines of this class are all intended for immediate use.

Decoction of Mallows.—Take of the roots of marsh-mallows, moderately dried, three ounces; raisins of the sun, one ounce; water, three pints.

Boil the ingredients in the water till one-third of it is consumed; afterwards strain the decoction, and let it stand for some time to settle. If the roots be thoroughly dried, they must be boiled till one-half of the water be consumed.

In coughs, and sharp defluxions upon the lungs, this decoction may be used for ordinary drink.

The Common Decoction.—Take of camomile-flowers, one ounce; elder-flowers, and sweet fennel-seeds, of each half an ounce; water, two quarts. Boil them for a little while, and then strain the decoction, or infuse the ingredients for some hours in boiling water.

This decoction is chiefly intended as the basis of clysters, to which other ingredients may be occasionally added. It will likewise serve as a common fomentation, spirit of wine or other things being added, in such quantity as the case may require.

Decoction of Logwood.—Boil three ounces of the shavings or chips of logwood, in four pints of water, till one half the liquor is wasted. Two or three ounces of simple cinnamon-water may be added to this decoction.

In fluxes of the belly, where the stronger astringents are improper, a tea-cupful of this decoction may be taken with advantage three or four times a-day.

Decoction of the Bark.—Boil an ounce of the Peruvian bark, grossly powdered, in a pint and a half of water to one pint; then strain the decoction. If a tea-spoonful of the weak spirit of vitriol be added to this medicine, it will render it both more agreeable and efficacious.

Compound Decoction of the Bark.—Take of Peruvian bark and Virginian snake-root, grossly powdered, each three drachms. Boil them in a pint of water to one-half. To the strained liquor add an ounce and a half of any aromatic water.

Sir John Pringle recommends this as a proper medicine towards the decline of malignant fevers, when the pulse is low, the voice weak, and the head affected with a stupor, but with little delirium. The dose is four table-spoonful every fourth or sixth hour.

Decoction of Sarsaparilla.—Take of fresh sarsaparilla-root, sliced and bruised, two ounces; shavings of guaiacum-wood, one ounce. Boil over a slow fire, in three quarts of water, to one; adding towards the end, half an ounce of sassafras-wood, and three drachms of liquorice. Strain the decoction.

This may either be employed as an assistant to a course of mercurial alteratives, or taken after the mercury has been used for some time. It strengthens the stomach, and restores flesh and vigour to habits emaciated by the venereal disease. It may also be taken in the rheumatism, and cutaneous disorders, proceeding from foulness of the blood and juices. For all these intentions it is greatly preferable to the *Decoction of Wood*. Dose: from a pint and a half to two quarts in the day.

The following decoction is said to be similar to that used by KENNEDY, in the cure of the venereal disease, and may supply the place of Lisbon diet drink:—

Take of sarsaparilla, three ounces; liquorice and mezereon-root, of each half an ounce; shavings of guaiacum and sassafras-wood, of each one ounce; crude antimony, powdered, an ounce and a half. Infuse these ingredients in eight pints of boiling water for twenty-four hours, then boil them till one half the water is consumed; afterwards strain the decoction. To be used in the same manner as the preceding.

Decoction of Seneka.—Take of Seneka or rattle-snake root, one ounce; water, a pint and a half. Boil to one pint, and strain.

This is recommended in the pleurisy, dropsy, rheumatism, and some obstinate disorders of the skin. The dose is two ounces, three or four times a-day or oftener, if the stomach will bear it.

White Decoction.—Take of the purest chalk, in powder, two ounces; gum arabic, half an ounce; water, three pints. Boil to one quart, and strain the decoction.

Proper drink in acute diseases, attended with, or inclining to, a looseness, and where acidities abound in the stomach or bowels. It is peculiarly proper for children when afflicted with sourness of the stomach, and for persons who are subject to the heart-burn. It may be sweetened with sugar, as it is used, and two or three ounces of simple cinnamon-water added to it.

An ounce of powdered chalk, mixed with two pints of water, will occasionally supply the place of this decoction, and also of the chalk julep.

DRAUGHTS.

THIS is a proper form for exhibiting such medicines as are intended to operate immediately, and which do not need to be frequently repeated, as purges, vomits, and

a few others, which are to be taken at one dose. Where a medicine requires to be used for any length of time, it is better to make up a large quantity of it at once, which saves both trouble and expense.

Anodyne Draught.—Take of liquid laudanum, twenty-five drops; simple cinnamon-water, an ounce; common syrup, two drachms. Mix them.

In excessive pain, where bleeding is not necessary, and in great restlessness, this composing draught may be taken and repeated occasionally.

Diuretic Draught.—Take of the diuretic salt, two scruples; syrup of poppies, two drachms; simple cinnamon-water, and common water, of each an ounce. In an obstruction or deficiency of urine.

Purging Draught.—Take of manna, an ounce; soluble tartar, or Rochel salt, from three to four drachms. Dissolve in three ounces of boiling water; to which add Jamaica pepper-water, half an ounce.

As manna sometimes will not sit upon the stomach, an ounce or ten drachms of the bitter purging salts, dissolved in four ounces of water, may be taken instead of the above.

Those who cannot take salts, may use the following draught:—Take of jalap in powder, a scruple; common water, an ounce; aromatic tincture, six drachms. Rub the jalap with twice its weight of sugar, and add to it the other ingredients.

Sweating Draught.—Take spirit of Mindererus, (*acetated solution of ammonia*) two ounces; salt of hartshorn, five grains; simple cinnamon-water, and syrup of poppies, of each half an ounce. Make them into a draught.

In recent colds and rheumatic complaints, this draught is of service. To promote its effects, however, the patient ought to drink freely of warm water-gruel, or of some other weak diluting liquor.

Vomiting Draughts.—Take of ipecacuanha, in powder, a scruple; water, an ounce; simple syrup, a drachm. Mix them. Persons who require a stronger vomit may add to this half a grain, or a grain, of emetic tartar.

Those who do not choose the powder, may take ten drachms of the ipecacuanha wine; or half an ounce of the wine, and an equal quantity of the syrup of squills.

ELECTUARIES.

ELECTUARIES are generally composed of the lighter powders mixed with syrup, honey, conserve, or mucilage, into such a consistence, that the powders may neither separate by keeping, nor the mass prove too stiff for swallowing. They receive chiefly the milder alterative medicines, and such as are not ungrateful to the palate.

Astringent electuaries, and such as have pulps of fruit in them, should be prepared only in small quantities; as astringent medicines lose their virtues by being kept in this form, and the pulps of fruits are apt to ferment.

For the extraction of pulps it will be necessary to boil unripe fruits, and ripe ones, if they are dried, in a small quantity of water, till they become soft. The pulp is then to be pressed out through a strong hair sieve, or thin cloth, and afterwards boiled to a due consistence, in an earthen vessel, over a gentle fire, taking care to prevent the matter from burning by continually stirring it. The pulps of fruit that are both ripe and fresh may be pressed out without any previous boiling.

Lenitive Electuary.—Take of senna, in fine powder, eight ounces; coriander-seed, also in powder, four ounces; pulp of tamarinds and of French prunes, each a pound. Mix the pulps and powders together, and with a sufficient quantity of simple syrup, reduce the whole into an electuary.

A tea-spoonful of this electuary, taken two or three times a-day, generally proves an agreeable laxative. It likewise serves as a convenient vehicle for exhibiting more active medicines, as jalap, scammony, and such like.

This may supply the place of the electuary of *Cassia*.

Electuary for the Dysentery.—Take of the Japonic confection, two ounces; Locatelli's balsam, one ounce; rhubarb in powder, half an ounce; syrup of marshmallows, enough to make an electuary.

About the bulk of a nutmeg should be taken twice or thrice a-day, as the symptoms and constitution may require.

It is often dangerous in dysenteries to give opiates and astringents, without interposing purgatives. The purgative is here joined with the above ingredients, which renders this a very safe and useful medicine for the purposes expressed in the title.

Electuary for the Epilepsy.—Take of Peruvian bark, in powder, an ounce; of powdered tin, and wild valerian root, each half an ounce; simple syrup, enough to make an electuary.

Dr. Mead directs a drachm of an electuary similar to this to be taken evening and morning, in the epilepsy, for the space of three months. It will be proper, however to discontinue the use of it for a few days every now and then. I have added the powdered tin, because the epilepsy often proceeds from worms.

Electuary for obstructed Menses.—Take of conserve of orange-peel, one ounce; steel-filings half an ounce; tartarized iron, two scruples; aromatic powder and powdered rhubarb, of each half a drachm; syrup, a sufficient quantity.

The bulk of a nutmeg of this taken every morning fasting, will be found an excellent remedy in obstructions of the menstrual evacuations.

Electuary of the Bark.—Take of Peruvian bark, in powder, three ounces; cascarilla, half an ounce; syrup of ginger, enough to make an electuary.

In the cure of obstinate intermitting fevers, the bark is assisted by the cascarilla. In hectic habits, however, it will be better to leave out the cascarilla, and put three drachms of crude sal ammoniac in its stead.

Electuary for the Piles.—Take flowers of sulphur, one ounce; cream of tartar, half an ounce; treacle, a sufficient quantity to form an electuary.

A tea-spoonful of this may be taken three or four times a-day.

Electuary for the Palsy.—Take of powdered mustard-seed, and conserve of roses each an ounce; syrup of ginger, enough to make an electuary.

A tea-spoonful of this may be taken three or four times a-day.

Electuary for the Rheumatism.—Take of conserve of roses, two ounces; cinnabar of antimony, levigated, an ounce and a half; gum guaiacum, in powder, an ounce; syrup of ginger, a sufficient quantity to make an electuary.

In obstinate rheumatisms, which are not accompanied with a fever, a tea-spoonful of this electuary may be taken twice a-day with considerable advantage.

EMULSIONS.

EMULSIONS, beside their use as medicines, are also proper vehicles for certain substances, which could not otherwise be conveniently taken in a liquid form. Thus camphor, triturated with almonds, readily unites with water into an emulsion. Pure oils, balsams, resins, and other similar substances, are likewise rendered miscible with water by the intervention of mucilages.

Common Emulsion.—Take of sweet almonds, an ounce; bitter almonds, a drachm; water, two pints.

Let the almonds be blanched, and beat up in a marble mortar; adding the water by little and little, so as to make an emulsion; afterwards let it be strained.

Arabic Emulsion.—This is made in the same manner as the above, adding to the almonds, while beating, two ounces and a half of the mucilage of gum arabic.

Where soft cooling liquors are necessary, these emulsions may be used as ordinary drink.

Camphorated Emulsion.—Take of Camphor, half a drachm; sweet almonds, half a dozen; white sugar, half an ounce; mint water, eight ounces. Grind the camphor and almonds well together in a stone mortar, and add by degrees the mint water; then strain the liquor, and dissolve in it the sugar.

In fevers, and other disorders which require the use of camphor, a table-spoonful of this emulsion may be taken every two or three hours.

Emulsion of Gum Ammoniac.—Take of gum ammoniac, two drachms; water, eight ounces. Grind the gum with the water poured upon it by little and little, till it is dissolved.

This emulsion is used for attenuating tough, viscid phlegm, and promoting expectoration. In obstinate coughs, two ounces of the syrup of poppies may be added to it. The dose is two table-spoonful three or four times a-day.

Oily Emulsion.—Take of soft water, six ounces; volatile aromatic spirit, two drachms; Florence oil, an ounce; shake them well together, and add, of simple syrup, half an ounce.

In recent colds and coughs, this emulsion is generally of service; but if the cough proves obstinate, it will succeed better when made with the paregoric elixir of the Edinburgh Dispensatory, instead of the volatile aromatic spirit. A table-spoonful of it may be taken every two or three hours.

EXTRACTS.

EXTRACTS are prepared by boiling the subject in water, and evaporating the strained

decoction to a due consistence. By this process some of the more active parts of plants are freed from the useless, indissoluble earthy matter, which makes the larger share of their bulk. Water, however, is not the only menstruum used in the preparation of extracts; sometimes it is joined with spirits, and at other times rectified spirit alone is employed for that purpose.

Extracts are prepared from a variety of different drugs, as the bark, gentian, jalap, &c.; but, as they require a troublesome and tedious operation, it will be more convenient for a private practitioner to purchase what he needs of them from a professed druggist, than to prepare them himself. Such of them as are generally used are inserted in our list of such drugs and medicines as are to be kept for private practice.

FOMENTATIONS.

FOMENTATIONS are generally intended either to ease pain, by taking off tension and spasm; or to brace and restore the tone and vigour of those parts to which they are applied. The first of these intentions may generally be answered by warm water, and the second by cold. Certain substances, however, are usually added to water with a view to heighten its effects, as anodynes, aromatics, astringents, &c. We shall therefore subjoin a few of the most useful medicated fomentations, that people may have it in their power to make use of them as they choose.

Anodyne Fomentation.—Take of white poppy-heads, two ounces; elder flowers, half an ounce; water, three pints. Boil till one pint is evaporated, and strain out the liquor.

This fomentation is used for relieving acute pain.

Aromatic Fomentation.—Take of Jamaica pepper, half an ounce; red wine, a pint. Boil them a little, and then strain the liquor.

This is intended, not only as a topical application for external complaints, but also for relieving the internal parts. Pains of the bowels, which accompany dysenteries and diarrheas, flatulent colics, uneasiness of the stomach, and retchings to vomit, are frequently abated by fomenting the abdomen and region of the stomach with the warm liquor.

Common Fomentation.—Take tops of wormwood and camomile flowers dried, of each two ounces; water, two quarts. After a slight boiling, pour off the liquor.

Brandy or spirit of wine may be added, in such quantity as the particular circumstances of the case shall require; but these are not always necessary.

Emollient Fomentation.—The same as the common decoction.

Strengthening Fomentation.—Take of oak bark, one ounce; granate peel, half an ounce; alum, two drachms; smith's forge-water, three pints. Boil the water with the bark and peel to the consumption of one-third; then strain the remaining decoction, and dissolve in it the alum.

This astringent liquor is employed as an external fomentation to weak parts; it may also be used internally.

GARGLES.

HOWEVER trifling this class of medicines may appear, they are by no means without their use. They seldom, indeed, cure diseases, but they often alleviate very disagreeable symptoms; as parchedness of the mouth, foulness of the tongue and fauces, &c.; they are peculiarly useful in fevers and sore throats. In the latter, a gargle will sometimes remove the disorder; and in the former, few things are more refreshing or agreeable to the patient, than to have his mouth frequently washed with some soft detergent gargle. One advantage of these medicines is, that they are easily prepared. A little barley-water and honey may be had any where; and if to these be added as much vinegar as will give them an agreeable sharpness, they will make a very useful gargle for softening and cleansing the mouth.

Attenuating Gargle.—Take of water, six ounces; honey, one ounce; nitre, a drachm and a half. Mix them.

This cooling gargle may be used either in the inflammatory quinsy, or in fevers, for cleaning the tongue and fauces.

Common Gargle.—Take of rose-water, six ounces; syrup of clove July-flowers, half an ounce; spirit of vitriol, a sufficient quantity to give it an agreeable sharpness. Mix them.

This gargle, besides cleansing the tongue and fauces, acts as a gentle repellent, and will sometimes remove a slight quinsy.

Detergent Gargle.—Take of the emollient gargle, a pint; tincture of myrrh, an ounce; honey, two ounces. Mix them.

When exulcerations require to be cleansed, or the excretion of tough viscid saliva promoted, this gargle will be of service.

Emollient Gargle.—Take an ounce of marshmallow roots, and two or three figs; boil them in a quart of water till near one-half of it be consumed; then strain out the liquor.

If an ounce of honey, and half an ounce of spirit of sal ammoniac, be added to the above, it will then be an exceedingly good *attenuating gargle*.

This gargle is beneficial in fevers, where the tongue and fauces are rough and parched, to soften these parts, and promote the discharge of saliva.

The learned and accurate Sir John Pringle observes, that in the inflammatory quinsy, or strangulation of the fauces, little benefit arises from the common gargles; that such as are of an acid nature do more harm than good, by contracting the emunctories of the saliva and mucus, and thickening those humours; that a decoction of figs in milk and water has a contrary effect, especially if some sal ammoniac be added; by which the saliva is made thinner, and the glands are brought to secrete more freely; a circumstance always conducive to the cure.

INFUSIONS.

VEGETABLES yield nearly the same properties to water by infusion as by decoction; and though they may require a longer time to give out their virtues in this way, yet it has several advantages over the other; since boiling is found to dissipate the finer parts of many bitter and aromatic substances, without more fully extracting their medicinal principles. Even from those vegetables which are weak in virtue, rich infusions may be obtained, by returning the liquor upon fresh quantities of the subject, the water loading itself more and more with the active parts; and these loaded infusions are applicable to valuable purposes in medicine, as they contain in a small compass the finer, more subtle, and active principles of vegetables, in a form readily miscible with the fluids of the human body.

Bitter Infusion.—Take tops of the lesser centaury and camomile flowers, of each half an ounce; yellow rind of lemon and orange-peel, carefully freed from the inner white part, of each two drachms. Cut them in small pieces, and infuse them in a quart of boiling water.

For indigestion, weakness of the stomach, or want of appetite, a tea-cupful of this infusion may be taken twice or thrice a-day.

Infusion of the Bark.—To an ounce of the bark, in powder, add four or five table-spoonsful of brandy, and a pint of boiling water. Let them infuse for two or three days.

This is one of the best preparations of the bark for weak stomachs. In disorders where the corroborating virtues of that medicine are required, a tea-cupful of it may be taken two or three times a-day.

Infusion of Carduus Benedictus (Blessed Thistle.)—Infuse an ounce of the dried leaves of carduus benedictus, or blessed thistle, in a pint of common water, for six hours, without heat; then filter the liquor through paper.

This light infusion may be given, with great benefit, in weakness of the stomach, where the common bitters do not agree. It may be flavoured at pleasure with cinnamon, or other aromatic materials.

Infusion of Linseed.—Take of linseed, two spoonsful; liquorice-root, sliced, half an ounce; boiling water, three pints. Let them stand to infuse by the fire for some hours, and then strain off the liquor.

If an ounce of the leaves of colt's-foot be added to these ingredients, it will then be the *Pectoral Infusion*. Both these are emollient mucilaginous liquors, and may be taken with advantage as ordinary drink in difficulty of making water; and in coughs and other complaints of the breast.

Infusion of Roses.—Take of red roses, dried, half an ounce; boiling water, a quart; vitriolic acid, commonly called oil of vitriol, half a drachm; loaf sugar, an ounce.

Infuse the roses in the water for four hours, in an unglazed earthen vessel; afterwards pour in the acid, and, having strained the liquor, add to it the sugar.

In an excessive flow of the menses, vomiting of blood, and other hæmorrhages, a tea-cupful of this gently astringent infusion may be taken every three or four hours. It likewise makes an exceedingly good gargle. As the quantity of roses used here can have little or no effect, an equally valuable medicine may be prepared by mixing the acid and water without infusion.

Infusion of Tamarinds and Senna.—Take of tamarinds, one ounce; senna, and crystals of tartar, each two drachms. Let these ingredients be infused four or five hours in a pint of boiling water; afterwards let the liquor be strained, and an ounce or

two of the aromatic tincture added to it. Persons who are easily purged may leave out either the tamarinds or the crystals of tartar.

This is an agreeable cooling purge, and supplies the place of the *Decoction of Tamarinds and Senna*. A tea-cupful may be given every half hour till it operates.

Spanish Infusion.—Take of Spanish juice, cut into small pieces, an ounce; salt of tartar, three drachms. Infuse in a quart of boiling water for a night. To the strained liquor add an ounce and a half of the syrup of poppies.

In recent colds, coughs, and obstructions of the breast, a tea-cupful of this infusion may be taken with advantage three or four times a-day.

Infusion for the Palsy.—Take of horse-radish root shaved, mustard-seed bruised, each four ounces; outer rind of orange-peel, one ounce. Infuse them in two quarts of boiling water, in a close vessel, for twenty-four hours.

In paralytic complaints, a tea-cupful of this warm stimulating medicine may be taken three or four times a-day. It excites the action of the solids, proves diuretic, and, if the patient be kept warm, promotes perspiration.

If two or three ounces of the dried leaves of marsh-trefoil be used instead of the mustard, it will make the *Antiscorbutic Infusion*.

JULEPS.

THE basis of juleps or draughts is generally common water, or some simple distilled water, with one-third or one-fourth its quantity of distilled spirituous water, and as much sugar or syrup as is sufficient to render the mixture agreeable. This is sharpened with vegetable or mineral acids, or impregnated with other medicines suitable to the intention.

Camphorated Julep.—Take of camphor, one drachm; rectified spirit of wine, ten drops; double-refined sugar, half an ounce; boiling distilled water, one pint. Rub the camphor first with the spirit of wine, then with the sugar; lastly, add the water by degrees, and strain the liquor.

In hysterical and other complaints where camphor is proper, this julep may be taken in the dose of a spoonful or two as often as the stomach will bear it.

Cordial Julep.—Take of simple cinnamon-water, four ounces; Jamaica pepper-water, two ounces; volatile aromatic spirit, and compound spirit of lavender, of each two drachms; syrup of orange-peel, an ounce. Mix them. Dose, two spoonful three or four times a-day, in disorders accompanied with great weakness and depression of spirits.

Expectorating Julep.—Take of the emulsion of gum ammoniac, six ounces; syrup of squills, two ounces. Mix them.

In coughs, asthmas, and obstructions of the breast, two table-spoonful of this julep may be taken every three or four hours.

Musk Julep.—Rub half a drachm of musk well together with half an ounce of sugar, and add to it, gradually, of simple cinnamon and peppermint-water, each two ounces; of the volatile aromatic spirit, two drachms.

In the low state of nervous fevers, hiccuping, convulsions, and other spasmodic affections, two table-spoonful of this julep may be taken every two or three hours.

Saline Julep.—Dissolve two drachms of salt of tartar in three ounces of fresh lemon juice, strained; when the effervescence is over, add of mint-water, and common water, each two ounces; of simple syrup, one ounce.

This removes sickness at the stomach, relieves vomiting, promotes perspiration, and may be of some service in fevers, especially of the inflammatory kind.

Vomiting Julep.—Dissolve four grains of emetic tartar in eight ounces of water, and add to it half an ounce of the syrup of clove July-flowers.

In the beginning of fevers, where there is no topical inflammation, this julep may be given in the dose of one table-spoonful every quarter of an hour till it operates. Antimonial vomits serve not only to evacuate the contents of the stomach, but likewise to promote the different excretions. Hence they are found in fevers to have nearly the same effects as *Dr. James's Powder*.

MIXTURES.

A MIXTURE differs from a julep in this respect, that it receives into its composition not only salts, extracts, and other substances dissoluble in water, but also earths, powders, and such substances as cannot be dissolved. A mixture is seldom either an elegant or agreeable medicine. It is nevertheless necessary. Many persons can take a mixture, who are not able to swallow a bolus or an electuary: besides, there are medicines which act better in this than in any other form.

Astringent Mixture.—Take simple cinnamon-water, and common water, of each three ounces; spirituous cinnamon-water, an ounce and a half; Japonic confection, half an ounce. Mix them.

In dysenteries which are not of long standing, after the necessary evacuations, a spoonful or two of this mixture may be taken every four hours, interposing every second or third day a dose of rhubarb.

The *Astringent Mixture*, which I have lately made use of with great success, is prepared thus:

Take powder of bole with opium, two drachms; cinnamon-water and penny-royal water, of each three ounces; spirituous cinnamon-water, six drachms; simple syrup, one ounce. Mix them, and take a table-spoonful four or five times a-day.

Diuretic Mixture.—Take of mint-water, five ounces; vinegar of squills, six drachms; sweet spirit of nitre, half an ounce; syrup of ginger, an ounce and a half. Mix them.

In obstructions of the urinary passages, two spoonfuls of this mixture may be taken twice or thrice a-day.

Laxative Absorbent Mixture.—Rub one drachm of magnesia alba in a mortar with ten or twelve grains of the best Turkey rhubarb, and add to them three ounces of common water; simple cinnamon-water, and syrup of sugar, of each one ounce.

As most diseases of infants are accompanied with acidities, this mixture may either be given with a view to correct these, or to open the body. A table-spoonful may be taken for a dose, and repeated three times a-day. To a very young child half a spoonful will be sufficient.

When the mixture is intended to purge, the dose may either be increased, or the quantity of rhubarb doubled.

This is one of the most generally useful medicines for children with which I am acquainted.

Saline Mixture.—Dissolve a drachm of the salt of tartar in four ounces of boiling water; and when cold, drop into it spirit of vitriol till the effervescence ceases; then add, of peppermint-water, two ounces; simple syrup, one ounce.

Where fresh lemons cannot be had, this mixture may occasionally supply the place of the saline julep.

Squill Mixture.—Take of simple cinnamon-water, five ounces; vinegar of squills, one ounce; syrup of marshmallows, an ounce and a half. Mix them.

This mixture, by promoting expectoration, and the secretion of urine, proves serviceable in asthmatic and dropsical habits. A table-spoonful of it may be taken frequently.

OINTMENTS, LINIMENTS, AND CERATES.

NOTWITHSTANDING the extravagant encomiums which have been best bestowed on different preparations of this kind, with regard to their efficacy in the cure of wounds, sores, &c. it is beyond a doubt, that the most proper application to a green wound is dry lint. But though ointments do not heal wounds and sores, yet they serve to defend them from the external air, and to retain such substances as may be necessary for drying, detaching, destroying proud flesh, and such like. For these purposes, however, it will be sufficient to insert only a few of the most simple forms, as ingredients of a more active nature can occasionally be added to them.

Yellow Basilicum Ointment.—Take of yellow wax, white resin, and frankincense, each a quarter of a pound; melt them together over a gentle fire; then add of hog's lard prepared, one pound. Strain the ointment while warm.

Employed for cleansing and healing wounds and ulcers.

Ointment of Calamine. (Turner's Cerate.)—Take of olive oil, a pint and a half; white wax, and calamine stone levigated, of each half a pound. Let the calamine stone, reduced into a fine powder, be rubbed with some part of the oil, and afterwards added to the rest of the oil and wax previously melted together, continually stirring them till quite cold.

An exceedingly good application in burns, and excoriations from whatever cause.

Emollient Ointment.—Take of palm oil, two pounds; olive oil, a pint and a half; yellow wax, half a pound; Venice turpentine, a quarter of a pound. Melt the wax in the oils over a gentle fire; then mix in the turpentine, and strain the ointment.

This supplies the place of *Althæa Ointment*. It may be used for anointing inflamed parts, &c.

Eye Ointment.—Take of hog's lard prepared, four ounces; white wax, two drachms; tatty prepared, one ounce; melt the wax with the lard over a gentle fire, and then sprinkle in the tatty, continually stirring them till the ointment is cold.

This preparation will be more efficacious, and of a better consistence, if two or three drachms of camphor be rubbed up with a little oil, and intimately mixed with it.

Another.—Take of camphor, and calamine stone levigated, each six drachms; verdegriese, well prepared, two drachms; hog's lard, and mutton-suet, prepared, of each two ounces. Rub the camphor well with the powder; afterwards mix in the lard and suet, continuing the triture till they be perfectly united.

This ointment has been long in esteem for diseases of the eyes. It ought, however, to be used with caution, when the eyes are much inflamed, or very tender.

Issue Ointment.—Mix half an ounce of Spanish flies, finely powdered, in six ounces of yellow basilicum ointment.

This ointment is chiefly intended for dressing blisters; in order to keep them open during pleasure.

Ointment of Lead.—Take of olive oil, half a pint; white wax, two ounces; sugar of lead, three drachms. Let the sugar of lead, reduced into a fine powder, be rubbed up with some part of the oil, and afterwards added to the other ingredients, previously melted together, continually stirring them till quite cold.

This cooling and gently astringent ointment may be used in all cases where the intention is to dry and skin over the part, as in scalding, &c.

Mercurial Ointment.—Take of quicksilver, two ounces; hog's lard, three ounces; mutton suet, one ounce. Rub the quicksilver with an ounce of the hog's lard in a warm mortar, till the globules be perfectly extinguished; then rub it up with the rest of the lard and suet, previously melted together.

The principal intention of this ointment is to convey mercury into the body by being rubbed upon the skin.

Ointment of Sulphur.—Take of hog's lard prepared, four ounces; flowers of sulphur, an ounce and a half; crude salammoniac, two drachms; essence of lemon, ten or twelve drops. Make them into an ointment.

This ointment, rubbed upon the parts affected, will generally cure the itch. It is both the safest and best application for that purpose, and, when made in this way, has no disagreeable smell.

Ointment for Diseases of the Skin.—Take of the ointment, commonly called *unguentum citrinum*, a drachm and a half; flour of brimstone and powder of hellebore, of each an ounce; hog's lard, three ounces; essence of lemon, or oil of thyme, from twenty to thirty drops, to correct the offensiveness of the smell. Make them into an ointment.

I have not only known many ordinary affections of the skin cured by this ointment, but even some of a very malignant nature, and approaching to leprosy.

White Ointment.—Take of olive-oil, one pint; white wax and spermaceti, of each three ounces. Melt them with a gentle heat, and keep them constantly and briskly stirring together, till quite cold.

If two drachms of camphor, previously rubbed with a small quantity of oil, be added to the above, it will make the *White camphorated Ointment*.

Liniment for Burns.—Take equal parts of Florence oil, or of fresh drawn linseed oil, and lime-water, shake them well together in a wide-mouthed bottle, so as to form a liniment.

This is found to be an exceedingly proper application for recent scalds or burns. It may either be spread upon a cloth, or the parts affected may be anointed with it twice or thrice a-day.

White Liniment.—This is made in the same manner as the white ointment, two-thirds of the wax being left out.

This liniment may be applied in cases of excoriation, where, on account of the largeness of the surface, the ointments with lead or calamine might be improper.

Liniment for the Piles.—Take of emollient ointment, two ounces; liquid laudanum, half an ounce. Mix these ingredients with the yolk of an egg, and work them well together.

Volatile Liniment.—Take of Florence oil, an ounce; spirit of hartshorn, half an ounce. Shake them together.

This liniment, made with equal parts of the spirit and oil, will be more efficacious, where the patient's skin is able to bear it.

Sir John Pringle observes, that in the inflammatory quinsey, a piece of flannel moistened with this liniment, and applied to the throat, to be renewed every four or five hours, is one of the most efficacious remedies; and that it seldom fails, after bleed-

hours, is one of the most efficacious remedies; and that it seldom fails, after bleeding, either to lessen or carry off the complaint. The truth of this observation I have often experienced.

Camphorated Oil.—Rub an ounce of camphor, with two ounces of Florence oil, in a mortar, till the camphor be entirely dissolved.

This antispasmodic liniment may be used in obstinate rheumatism, and in some other cases accompanied with extreme pain and tension of the parts.

PILLS.

MEDICINES which operate in a small dose, and whose disagreeable taste or smell makes it necessary that they should be concealed from the palate, are more commodiously exhibited in this form. No medicine, however, that is intended to operate quickly, ought to be made into pills, as they often lie for a considerable time on the stomach before they are dissolved, so as to produce any effect.

As the ingredients which enter the composition of pills are generally so contrived, that one pill of an ordinary size may contain about five grains of the compound, in mentioning the dose we shall only specify the number of pills to be taken: as one, two, three, &c.

Composing Pill.—Take of purified opium, ten grains; Castile soap, half a drachm. Beat them together, and form the whole into twenty pills.

When a quieting draught will not sit upon the stomach, one, two, or three of these pills may be taken, as occasion requires.

Deobstruent Pill.—Take salt of steel; succotrine aloes; myrrh in powder; of each a drachm. Make into forty pills, of which two are to be taken evening and morning.

I have found these pills of excellent service in obstructions of the *menes*. The late Dr. WATKINSON made it his dying request, that I would insert this prescription in the "Domestic Medicine," which he said would be immortal, and that "his soul panted for immortality."

Fatid Pill.—Take of asafetida, half an ounce; simple syrup, as much as is necessary to form it into pills.

In hysteric complaints, four or five pills, of an ordinary size, may be taken twice or thrice a-day. They may likewise be of service to persons afflicted with the asthma.

When it is necessary to keep the body open, a proper quantity of rhubarb, aloes, or jalap, may occasionally be added to the above mass.

Hemlock Pill.—Take any quantity of the extract of hemlock, and adding to it about a fifth part its weight of the powder of the dried leaves, form it into pills of the ordinary size.

The extract of hemlock may be taken from one grain to several drachms in the day. The best method, however, of using these pills, is to begin with one or two, and to increase the dose gradually, as far as the patient can bear them, without any remarkable degree of stupor or giddiness.

Mercurial Pill.—Take of purified quicksilver and honey, each half an ounce. Rub them together in a mortar, till the globules of mercury are perfectly extinguished; then add, of Castile soap, two drachms; powdered liquorice, or crumb of bread, a sufficient quantity to give the mass a proper consistence for pills.

When stronger mercurial pills are wanted, the quantity of quicksilver may be doubled.

The dose of these pills is different, according to the intention with which they are given. As an alterant, two or three may be taken daily. To raise a salivation, four or five will be necessary.

Equal parts of the above pill and powdered rhubarb, made into a mass, with a sufficient quantity of simple syrup, will make a *Mercurial Purging Pill*.

Mercurial Sublimate Pill.—Dissolve fifteen grains of the corrosive sublimate of mercury in two drachms or the saturated solution of crude sal ammoniac, and make it into a paste, in a glass mortar, with a sufficient quantity of the crumb of bread. This must be formed into one hundred and twenty pills.

For the venereal disease, four of these pills may be taken twice a-day, as an alterant three, and for worms, two.

This pill, which is the most agreeable form of exhibiting the sublimate, has been found efficacious, not only in curing the venereal disease, but also in killing and expelling worms, after other powerful medicines had failed.*

Plummer's Pill.—Take of calomel, or sweet mercury, and precipitated sulphur of

* See a paper on this subject in the Edinburgh Physical and Literary Essays, by the ingenious Dr. John Gardener.

antimony, each three drachms; extract of liquorice, two drachms. Rub the sulphur and mercury well together: afterwards add the extract, and with a sufficient quantity of the mucilage of gumarabic, make them into pills.

This pill has been found a powerful, yet safe alterative in obstinate cutaneous disorders; and has completed a cure after salivation had failed. In venereal cases it has likewise produced excellent effects. Two or three pills of an ordinary size may be taken night and morning, the patient keeping moderately warm, and drinking after each dose a draught of decoction of the woods, or of sarsaparilla.

Purging Pill.—Take of succotrine aloes, and Castile soap, each two drachms; of simple syrup, a sufficient quantity to make them into pills.

Four or five of these pills will generally prove a sufficient purge. For keeping the body gently open, one may be taken night and morning. They are reckoned both deobstruent and stomachic, and will be found to answer all the purposes of Dr. Anderson's pills, the principal ingredient of which is aloes.

Where aloetic purges are improper, the following pills may be used:—

Take extract of jalap, and vitriolated tartar, of each two drachms; syrup of ginger, as much as will make them of a proper consistence for pills. To be taken in the same quantity as the above.

Purgative Pill.—Take powder of succotrine aloes, one drachm; of gum sagapene in powder, half a drachm; of gamboge, and gum arabic in powder, each, one scruple; essential oil of camomile, ten drops; syrup of buckthorn, a sufficient quantity; beat the whole into a mass, and divide into thirty-two pills.

This pill was contrived by that eminent physician, the late Dr. GEORGE FORDYCE. It is an excellent purgative, where the bowels are torpid, as in paralytic affections. One or two may be taken at bed-time. A. P. B.

Pill for the Bile.—Take gum pill and colocynth pill, each a drachm. Beat them together, and make the mass into thirty pills.

In bilious and nervous patients, where it was necessary to keep the body gently open, I have found these pills answer the purpose extremely well. I generally give one over-night, and another next morning, once or twice a-week. But the dose must be regulated by the effect.

Pill for the Jaundice.—Take of Castile soap, succotrine aloes, and rhubarb, of each one drachm. Make them into pills, with a sufficient quantity of syrup or mucilage.

These pills, as their title expresses, are chiefly intended for the jaundice, which, with the assistance of proper diet, they will often cure. Five or six of them may be taken twice a-day, more or less, as is necessary to keep the body open. It will be proper, however, during their use, to interpose now and then a vomit of ipecacuanha or tartar emetic.

Stomachic Pill.—Take extract of gentian, two drachms; powdered rhubarb, and vitriolated tartar, of each one drachm; oil of mint, thirty drops; simple syrup, a sufficient quantity.

Three or four of these pills may be taken twice a-day, for invigorating the stomach, and keeping the body gently open.

Squill Pills.—Take powder of dried squills, a drachm and a half; gum ammoniac, and cardamom seeds, in powder, of each three drachms; simple syrup, a sufficient quantity.

In dropsical and asthmatic complaints, two or three of these pills may be taken twice a-day, or oftener, if the stomach will bear them.

Strengthening Pills.—Take soft extract of the bark, and salt of steel, each a drachm. Make into pills.

In disorders arising from excessive debility, or relaxation of the solids, as the *chlorosis*, or green sickness, two of these pills may be taken three times a-day.

PLASTERS.

PLASTERS ought to be of a different consistence, according to the purposes for which they are intended. Such as are to be applied to the breasts or stomach, ought to be soft and yielding; while those designed for the limbs should be firm and adhesive.—It has been supposed, that plasters might be impregnated with the virtues of different vegetables, by boiling the recent vegetable with the oil employed for the composition of the plaster; but this treatment does not communicate to the oils any valuable qualities.—The *calces* of lead boiled with oils unite with them into a plaster of a proper consistence, which makes the basis of several other plasters. In boiling these compositions, a quantity of hot water must be added from time to time to prevent the plaster from burning or growing black. This, however, should be done with care, lest it cause the matter to explode.

Common Plaster.—Take of common olive oil, six pints; litharge, reduced to a fine powder, two pounds and a half. Boil the litharge and oil together over a gentle fire, continually stirring them, and keeping always about a half a gallon of water in the vessel; after they have boiled about three hours, a little of the plaster may be taken out and put into cold water, to try if it be of a proper consistence: when that is the case, the whole may be suffered to cool, and the water well pressed out of it with the hands.

This plaster is generally applied in slight wounds and excoriations of the skin. It keeps the part soft and warm, and defends it from the air, which is all that is necessary in such cases. Its principal use, however, is to serve as a basis for other plasters.

Adhesive Plaster.—Take of common plaster, half a pound; of Burgundy pitch, a quarter of a pound. Melt them together.

This plaster is principally used for keeping on other dressings.

Anodyne Plaster.—Melt an ounce of adhesive plaster, and when it is cooling, mix with it a drachm of powdered opium, and the same quantity of camphor, previously rubbed up with a little oil.

This plaster generally gives ease in acute pains, especially of the nervous kind.

Blistering Plaster.—Take of Venice turpentine, six ounces; yellow wax, two ounces; Spanish flies in fine powder, three ounces; powdered mustard, one ounce. Melt the wax, and while it is warm, add to it the turpentine, taking care not to evaporate it by too much heat. After the turpentine and wax are sufficiently incorporated, sprinkle in the powder, continually stirring the mass till it be cold.

Though this plaster is made in a variety of ways, one seldom meets with it of a proper consistence. When compounded with oils and other greasy substances, its effects are blunted, and it is apt to run; while pitch and resin render it too hard and very inconvenient.

When the blistering-plaster is not at hand, its place may be supplied by mixing with any soft ointment a sufficient quantity of powdered flies; or by forming them into a paste with flour and vinegar.

Blistering-plasters prove highly disagreeable to many people, by occasioning strangury. I have, therefore, of late used a plaster, in which a small quantity of blistering-salve has been mixed with the Burgundy pitch-plaster. I lay it over the part affected, and suffer it to remain as long as it will stick. The blistering-plaster loses its effect in a few hours, whereas this will act for many days, or even weeks, and seldom fails to remove pain, or slight obstructions.

Gum Plaster.—Take of the common plaster, four pounds; gum ammoniac and galbanum, strained, of each half a pound. Melt them together, and add, of Venice turpentine, six ounces.

This plaster is used as a digestive, and likewise for discussing indolent tumours.

Mercurial Plaster.—Take of common plaster, one pound; of gum ammoniac, strained, half a pound. Melt them together, and, when cooling, add eight ounces of quicksilver, previously extinguished by triture, with three ounces of hog's lard.

This plaster is recommended in pains of the limbs arising from a venereal cause. Indurations of the glands, and other indolent tumours, are likewise found sometimes to yield to it.

Stomach Plaster.—Take of gum plaster, half a pound; camphorated oil, an ounce and a half; black pepper, or capsicum, where it can be had, one ounce. Melt the plaster, and mix with it the oil; then sprinkle in the pepper, previously reduced to a fine powder.

An ounce or two of this plaster, spread upon soft leather, and applied to the region of the stomach, will be of service in flatulencies arising from hysteric and hypochondriac affections. A little of the expressed oil of mace, or a few drops of the essential oil of mint, may be rubbed upon it before it is applied.

This may supply the place of the *Anti-hysteric Plaster*.

Warm Plaster.—Take of gum plaster, one ounce; blistering-plaster, two drachms. Melt them together over a gentle fire.

This plaster is useful in the sciatica and other fixed pains of the rheumatic kind: it ought, however, to be worn for some time, and to be renewed at least once a-week. If this is found to blister the part, which is sometimes the case, it must be made with a smaller proportion of the blistering-plaster.

Wax Plaster.—Take of yellow wax, one pound; white resin, half a pound; muton-suet, three quarters of a pound. Melt them together.

This is generally used instead of the *Mellot Plaster*. It is a proper application after blisters, and in other cases where a gentle digestive is necessary.

POULTICES.

THROUGH some oversight, this article was omitted in the earlier editions, though it relates to a class of medicines by no means unimportant. Poultrices are often beneficial, even in the most simple form; but more so, when employed to retain more active medicines,—to keep them in contact with the skin,—and to fit it for their absorption. Every nurse knows how to make a poultice.

A poor woman who had received a very dangerous wound in the tendons of her thumb from a rusty nail, called upon me some little time since. As her case properly belonged to the department of surgery, I advised her to apply to the hospital; but the official hirelings there refused to take her in, though I always understood that they were *obliged to take in accidents*. It seems, however, that some very confined meaning was annexed to this word by the surgeon on duty, and that he did not think the danger of a locked jaw to be an *accident* as deserving of his pity and immediate assistance, as a broken arm, or dislocated ancle. The poor woman came back to me; and, as her situation became every moment more and more alarming, the pain and inflammation having reached as high as the arm-pit, I advised her to apply to the whole hand and arm a large poultice, with an ounce of laudanum sprinkled over it, and to renew the poultice twice a-day. This she did with so much success, that the thumb is now quite well, though the accident did not happen above three weeks ago.

Alarming as the case was, I had some reason to rely on the efficacy of the poultice, from a former trial somewhat similar. One of those girls, who are employed by bookbinders in stitching the sheets, having wounded her finger with the three-edged needle used on such occasions, soon felt the pain shoot upwards with deadly tendency. I ordered her to apply the same sort of poultice with laudanum, which had the same happy effect.

Both these patients made use of the *Common Poultice*; but I prefer one made of linseed flour, which is more easily prepared, and keeps moist longer than any other.

POWDERS.

THIS is one of the most simple forms in which medicine can be administered. Many medicinal substances, however, cannot be reduced into powder, and others are too disagreeable to be taken in this form. The lighter powders may be mixed in any agreeable thin liquor, as tea or water-gruel. The more ponderous will require a more consistent vehicle, as syrup, conserve, jelly, or honey. Gums, and other substances, which are difficult to powder, should be pounded along with the drier ones; but those which are too dry, especially aromatics, ought to be sprinkled during their pulverization with a few drops of any proper water. Aromatic powders are to be prepared only in small quantities at a time, and kept in glass vessels closely stopped. Indeed, no powders ought to be exposed to the air, or kept too long, otherwise their virtues will be in great measure destroyed.

Astringent Powder.—Take of alum and Japan earth, each two drachms. Pound them together, and divide the whole into ten or twelve doses.

In an immoderate flow of the *menses*, and other hæmorrhages, one of these powders may be taken every hour, or every half-hour, if the discharge be violent.

Powder of Bole.—Take of Bole armenic, or French Bole, two ounces; cinnamon, one ounce; tormentil root and gum arabic of each six drachms; long pepper, one drachm. Let all these ingredients be reduced into a powder.

This warm, glutinous, astringent powder is given in fluxes, and other disorders where medicines of that class are necessary, in the dose of a scruple, or half a drachm.

If a drachm of opium be added, it will make the *Powder of Bole with Opium*, which is a medicine of considerable efficacy. It may be taken in the same quantity as the former, but not above twice or thrice a-day.

Carminative Powder.—Take of coriander-seed, half an ounce; ginger, one drachm; nutmegs, half a drachm; fine sugar, a drachm and a half. Reduce them into powder for twelve doses.

This powder is employed for expelling flatulencies arising from indigestion, particularly those to which hysteric and hypochondriac persons are so liable. It may likewise be given in small quantities to children in their food, when troubled with gripes.

Diuretic Powder.—Take of gum arabic four ounces; purified nitre, one ounce. Pound them together, and divide the whole into twenty-four doses.

During the first stage of the venereal disease, one of these cooling powders may be taken three times a-day, with considerable advantage.

Aromatic Opening Powder.—Take the best Turkey rhubarb, cinnamon, and fine sugar, each two drachms. Let the ingredients be pounded, and afterwards mixed well together.

When flatulency is accompanied by costiveness, a tea-spoonful of this powder may be taken once or twice a-day, according to circumstances.

Saline Laxative Powder.—Take of soluble tartar, and cream of tartar, each one drachm; purified nitre, half a drachm. Make them into a powder.

In fevers and other inflammatory disorders, where it is necessary to keep the body gently open, one of these cooling laxative powders may be taken in a little gruel, and repeated occasionally.

Steel Powder.—Take filings of steel, and loaf sugar, of each two ounces; ginger, two drachms. Pound them together.

In obstructions of the *menses*, and other cases where steel is proper, a tea-spoonful of this powder may be taken twice a-day, and washed down with a little wine and water.

Sudorific Powder.—Take purified nitre and vitriolated tartar, of each half an ounce; opium and ipecacuanha, of each one drachm. Mix the ingredients, and reduce them to a fine powder.

This is generally known by the name of *Doner's Powder*. It is a powerful sudorific. In obstinate rheumatisms, and other cases where it is necessary to excite a copious sweat, this powder may be administered in the dose of a scruple or half a drachm. Some patients will require two scruples. It ought to be accompanied with the plentiful use of some warm diluting liquor.

Worm Powder.—Take of tin reduced into a fine powder, an ounce; *Aethiops mineral*, two drachms. Mix them well together, and divide the whole into six doses.

One of these powders may be taken in a little syrup, honey, or treacle, twice a-day. After they have been all used, the following anthelmintic purge may be proper.

Purging Worm Powder.—Take the powdered rhubarb, a scruple; scammony and calomel of each five grains. Rub them together in a mortar for one dose. For children, however, these doses must be lessened according to their age.

If the powder of tin be given alone, its dose may be considerably increased. The late Dr. Alston gave it to the amount of two ounces in three days, and says, when thus administered, that it proves an egregious anthelmintic. He purged his patients both before they took the powder and afterwards.

Powder for the Tape Worm.—Early in the morning the patient is to take in any liquid two or three drachms, according to his age and constitution, of the root of the male fern reduced into a fine powder. About two hours afterwards, he is to take of calomel and resin of scammony, each ten grains; gum gamboge, six grains. These ingredients must be finely powdered and given in a little syrup, honey, treacle, or any thing that is most agreeable to the patient. He is then to walk gently about, now and then drinking a dish of weak green tea, till the worm is passed. If the powder of the fern produces nausea or sickness, it may be removed by sucking the juice of an orange or lemon.

This medicine, which had been long kept a secret abroad for the cure of the tape-worm, was some time ago purchased by the French king, and made public for the benefit of mankind. Not having had an opportunity of trying it, I can say nothing from experience concerning its efficacy. It seems, however, from its ingredients, to be an active medicine, and ought to be taken with care. The dose here prescribed is sufficient for the strongest patient; it must, therefore, be reduced according to the age and constitution.

SYRUPS.

SYRUPS were some time ago looked upon as medicines of considerable value. They are at present, however, regarded chiefly as vehicles for medicines of greater efficacy, and are used for sweetening draughts, juleps, or mixtures; and for reducing the lighter powders into boluses, pills, and electuaries. As all these purposes may be answered by the simple syrup alone, there is little occasion for any other; especially as they are seldom found but in a state of fermentation; and as the dose of any medicine given in this form is very uncertain. Persons who serve the public must keep whatever their customers call for; but, to the private practitioner, nine-tenths of the syrups usually kept in the shops are unnecessary.

Simple Syrup.—Is made by dissolving in water, either with or without heat, about double its weight of fine sugar.

If twenty-five drops of laudanum be added to an ounce of the simple syrup, it will supply the place of diacodium, or the syrup of poppies, and will be found a more safe and certain medicine.

The lubricating virtues of the syrup of marshmallows may likewise be supplied, by adding to the common syrup a sufficient quantity of mucilage of gum arabic.

Those who choose to preserve the juice of lemons in form of syrup, may dissolve in it, by the heat of a warm bath, nearly double its weight of fine sugar. The juice ought to be previously strained, and suffered to stand till it settles.

The syrup of ginger is sometimes of use as a warm vehicle for giving medicine to persons afflicted with flatulency. It may be made by infusing two ounces of bruised ginger in two pints of boiling water for twenty-four hours. After the liquor has been strained, and has stood to settle for some time, it may be poured off, and a little more than double its weight of fine powdered sugar dissolved in it.

TINCTURES, ELIXIRS, &c.

RECTIFIED spirit is the direct menstruum of the resins and essential oils of vegetables, and totally extracts these active principles from sundry substances, which yield them to water, either not at all, or only in part.

It dissolves likewise those parts of animal substances in which their peculiar smells and tastes reside. Hence, the tinctures prepared with rectified spirits form an useful and elegant class of medicines, possessing many of the most essential virtues of simples, without being clogged with their inert or useless parts.

Water, however, being the proper menstruum of the gummy, saline, and saccharine parts of medicinal substances, it will be necessary, in the preparation of several tinctures, to make use of a weak spirit, or a composition of rectified spirit and water.

Aromatic Tincture.—Infuse two ounces of Jamaica pepper in two pints of brandy, without heat, for a few days: then strain off the tincture.

This simple tincture will sufficiently answer all the intentions of the more costly preparations of this kind. It is rather too hot to be taken by itself; but is very proper for mixing with such medicines as might otherwise prove too cold for the stomach.

Compound Tincture of the Bark.—Take of Peruvian bark, two ounces; Seville orange-peel and cinnamon, of each half an ounce. Let the bark be powdered and the other ingredients bruised: then infuse the whole in a pint and a half of brandy, for five or six days, in a close vessel; afterwards strain off the tincture.

The dose is from one drachm to three or four, every fifth or sixth hour. It may be given in any suitable liquor, and occasionally sharpened with a few drops of the spirits of vitriol.

This tincture is not only beneficial in intermitting fevers, but also in the slow, nervous, and putrid kinds, especially towards their decline.

Volatile Fetid Tincture.—Infuse two ounces of asafoetida in one pint of volatile aromatic spirit, for eight days, in a close bottle, frequently shaking it; then strain the tincture.

This medicine is beneficial in hysteric disorders, especially when attended with lowness of spirits, and faintings. A tea-spoonful of it may be taken in a glass of wine, or a cup of penny-royal tea.

Volatile Tincture of Gum Guaiacum.—Take of gum guaiacum, four ounces; volatile aromatic spirit, a pint. Infuse without heat, in a vessel well stopped, for a few days; then strain off the tincture.*

In rheumatic complaints, a tea-spoonful of this tincture may be taken in a cup of the infusion of water trefoil, twice or thrice a-day.

Tincture of Black Hellebore.—Infuse two ounces of the roots of black hellebore, bruised, in a pint of proof spirit, for seven or eight days; then filter the tincture through paper. A scruple of cochineal may be infused along with the roots, to give the tincture a colour.

In obstructions of the menses, a tea-spoonful of this tincture may be taken in a cup of camomile or penny-royal tea twice a-day.

Astringent Tincture.—Digest two ounces of gum kino, in a pint and a half of brandy, for eight days; afterwards strain it for use.

This tincture, though not generally known, is a good astringent medicine. With this view, an ounce, or more, of it may be taken three or four times a-day.

Tincture of Myrrh and Aloes.—Take of gum myrrh, an ounce and a half; hepatic aloes, one ounce. Let them be reduced to a powder, and infused in two pints of rectified spirits, for six days, in a gentle heat; then strain the tincture.

This is principally used by surgeons for cleansing foul ulcers, and restraining the progress of gangrenes. It is also, by some, recommended as a proper application to green wounds.

* A very good tincture of guaiacum, for domestic use, may be made by infusing two or three ounces of the gum in a bottle of rum or brandy.

*Tincture of Opium (Liquid Laudanum).—*Take of crude opium, two ounces; spirituous aromatic water, and mountain wine, of each ten ounces. Dissolve the opium, sliced, in the wine, with a gentle heat, frequently stirring it; afterwards add the spirit, and strain off the tincture.

As twenty-five drops of this tincture contain about a grain of opium, the common dose may be from twenty to thirty drops.

*Sacred Tincture, or Tincture of Hiera Picra.—*Take of succotrine aloes in powder, one ounce; Virginian snake-root and ginger, of each two drachms. Infuse in a pint of mountain wine, and half a pint of brandy, for a week, frequently shaking the bottle; then strain off the tincture.

This is a safe and useful purge for persons of a languid and phlegmatic habit: but it is thought to have better effects, taken in small doses as a laxative.

The dose, as a purge, is from one to two ounces.

*Compound Tincture of Senna.—*Take of senna, one ounce; jalap, coriander-seeds, and cream of tartar, of each half an ounce. Infuse them in a pint and a half of French brandy for a week; then strain the tincture, and add to it four ounces of fine sugar.

This is an agreeable purge, and answers all the purposes of the *Elixir Salutis*, and of *Daffy's Elixir*. The dose is from one to two or three ounces.

*Tincture of Spanish Flies.—*Take of Spanish flies, reduced to a fine powder, two ounces; spirit of wine, one pint. Infuse for two or three days; then strain off the tincture.

This is intended as an acrid stimulant for external use. Parts affected with the palsy, or chronic rheumatism, may be frequently rubbed with it.

*Tincture of the Balsam of Tolu.—*Take of the balsam of Tolu, an ounce and a half; rectified spirit of wine, a pint. Infuse in a gentle heat until the balsam is dissolved; then strain the tincture.

This tincture possesses all the virtues of the balsam. In coughs, and other complaints of the breast, a tea-spoonful or two of it may be taken on a bit of loaf-sugar. But the best way of using it is in syrup. An ounce of the tincture properly mixed with two pounds of simple syrup, will make what is commonly called the *Balsamic Syrup*.

*Tincture of Rhubarb.—*Take of rhubarb, two ounces and a half; lesser cardamom-seeds half an ounce; brandy, two pints. Digest for a week, and strain the tincture.

Those who choose to have a vinous tincture of rhubarb may infuse the above ingredients in a bottle of Lisbon wine, adding to it about two ounces of proof spirits.

If half an ounce of gentian root, and a drachm of Virginian snake-root, be added to the above ingredients, it will make the bitter tincture of rhubarb.

All these tinctures are designed as stomachics and corroborants as well as purgatives. In weakness of the stomach, indigestion, laxity of the intestines, fluxes, colicky and such like complaints, they are frequently of great service. The dose is from half a spoonful to three or four spoonfuls or more, according to the circumstances of the patient, and the purposes it is intended to answer.

*The Tonic Tincture.—*Mix two ounces of the compound tincture of Peruvian bark with the like quantity of the volatile tincture of Valerian; and of this mixture a tea-spoonful in a glass of wine or water is to be taken three or four times a-day.

I have long made use of this tincture for the relief of those peculiar affections of the stomach and bowels, such as indigestion, &c. which generally accompany nervous diseases. I do not say that the tincture will cure those complaints, nor do I know of any medicine that will; but where a complete cure cannot be rationally expected, relief is certainly a very desirable object.

*Paregoric Elixir.—*Take of flowers of benzoin, half an ounce; opium, two drachms. Infuse in one pound of the volatile aromatic spirit, for four or five days, frequently shaking the bottle; afterwards strain the elixir.

This is an agreeable and safe way of administering opium. It eases pain, allays tickling coughs, relieves difficult breathing, and is useful in many disorders of children, particularly the whooping-cough. The dose to an adult is from fifty to an hundred drops.

*Sacred Elixir.—*Take of rhubarb, cut small, ten drachms; succotrine aloes, in powder, six drachms; lesser cardamom-seeds, half an ounce; French brandy, two pints. Infuse for two or three days, and then strain the elixir.

This useful stomachic purge may be taken from one ounce to an ounce and a half.

*Stomachic Elixir.—*Take of gentian root, two ounces; Curassao oranges, one ounce; Virginian snake-root, half an ounce. Let the ingredients be bruised, and infused for three or four days in two pints of French brandy; afterwards strain out the elixir.

This is an excellent stomach bitter. In flatulencies, indigestion, want of appetite,

and such like complaints, a small glass of it may be taken twice a-day. It likewise relieves the gout in the stomach, when taken in a large dose.

Acid Elixir of Vitriol.—Take of the aromatic tincture, one pint; oil of vitriol, three ounces. Mix them gradually, and after the fæces have subsided, filter the elixir through paper, in a glass funnel.

This is one of the best medicines which I know for hysteric and hypochondriac patients, afflicted with flatulencies arising from relaxation or debility of the stomach and intestines. It will succeed where the most celebrated stomachic bitters have no effect. The dose is from ten to forty drops, in a glass of wine or water, or a cup of any bitter infusion, twice or thrice a-day. It should be taken when the stomach is most empty.

Camphorated Spirit of wine.—Dissolve an ounce of camphor in a pint of rectified spirits.

This solution is chiefly employed as an embrocation in bruises, palsies, the chronic rheumatism, and for preventing gangrenes. The above quantity of camphor, dissolved in half a pound of the volatile aromatic spirit, makes *Ward's Essence*.

Spirit of Mindererus. (*Solution of acetated Ammonia.*) Take of volatile sal ammoniac, any quantity. Pour on it gradually distilled vinegar, till the effervescence ceases.

This medicine is useful in promoting a discharge both by the skin and urinary passage. It is also a good external application in strains and bruises. When intended to raise a sweat, half an ounce of it in a cup of warm gruel may be given to the patient in bed every hour till it has the desired effect.

VINEGARS.

VINEGAR is an acid produced from vinous liquors by a second fermentation. It is an useful medicine both in inflammatory and putrid disorders. Its effects are, to cool the blood, quench thirst, counteract a tendency to putrefaction, and allay inordinate motions of the system. It likewise promotes the natural secretions, and in some cases excites a copious sweat, where the warm medicines, called alexipharmic, tend rather to prevent that salutary evacuation.

Vinegar is also advantageously employed in burns and scalds; by keeping the parts constantly wet with it, by means of linen rags. "In severe burns and scalds," observes Mr. Cleghorn, "which have recently happened, and which are attended with large blisters, excoriations or loss of substance, the vinegar must be constantly applied till the heat and pain nearly cease, which will happen in from two to eight hours, according as the injury is more or less severe. The sores must be covered with rags or cloths well wetted, which, as often as they dry, or any sensation of pain or heat returns, must be wetted afresh with the vinegar, for two, three, or four hours."

Weakness, faintings, vomitings, and other hysteric affections, are often relieved by vinegar applied to the mouth and nose, or received into the stomach. It is of excellent use also in correcting many poisonous substances, when taken into the stomach; and in promoting their expulsion, by the different emunctories, when received into the blood.

Vinegar is not only an useful medicine, but serves likewise to extract, in tolerable perfection, the virtues of several other medicinal substances. Most of the odoriferous flowers impart to it their fragrance, together with a beautiful purplish or red colour. It also assists or coincides with the intention of squills, garlic, gum ammoniac, and several other valuable medicines.

These effects, however, are not to be expected from every thing that is sold under the name of vinegar, but from such as is sound and well prepared.

The best vinegars are those prepared from French wines.

It is necessary for some purposes that the vinegar be distilled; but as this operation requires a particular chemical apparatus, we shall not insert it.

Vinegar of Litharge.—Take of litharge, half a pound; strong vinegar, two pints. Infuse them together in a moderate heat for three days, frequently shaking the vessel; then filter the liquor for use.

This medicine is little used, from a general notion of its being dangerous. There is reason, however, to believe, that the preparations of lead with vinegar are possessed of some valuable properties, and that they may be used in many cases with safety and success.

A preparation of a similar nature with the above has of late been extolled by Goulard, a French surgeon, as a safe and extensively-useful medicine, which he calls the *Extract of Saturn*, and orders to be made in the following manner:—

Take of litharge, one pound; vinegar made of French wine, two pints. Put them together into a glazed earthen pipkin, and let them boil, or rather simmer, for an hour, or an hour and a quarter, taking care to stir them all the while with a wooden

spatula. After the whole has stood to settle, pour off the liquor which is upon the top into bottles for use.

With this extract Goulard makes his *vegeto mineral water*,* which he recommends in a great variety of external disorders, as inflammations, burns, bruises, sprains, ulcers, &c.

He likewise prepares with it a number of other forms of medicine, as poultices, plasters, ointments, powders, &c.

Vinegar of Roses.—Take of red roses, half a pound; strong vinegar, half a gallon. Infuse in a close vessel for several weeks, in a gentle heat; and then strain off the liquor.

This is principally used as an embrocation for headachs, &c.

Vinegar of Squills.—Take of dried squills, two ounces; distilled vinegar, two pints. Infuse for ten days or a fortnight in a gentle degree of heat, afterwards strain off the liquor, and add to it about a twelfth part of its quantity of proof spirits.

This medicine has good effects in disorders of the breast, occasioned by a load of viscid phlegm. It is also of use in hydropic cases for promoting a discharge of urine.

The dose is from two drachms to two ounces, according to the intention for which it is given. When intended to act as a vomit, the dose ought to be large. In other cases it must not only be exhibited in small doses, but also mixed with cinnamon-water, or some other agreeable aromatic liquor, to prevent the nausea it might otherwise occasion.

WATERS BY INFUSION, &c.

LIME WATER.—Pour two gallons of water gradually upon a pound of fresh burnt quicklime; and when the ebullition ceases, stir them well together; then suffer the whole to stand at rest, that the lime may settle, and afterwards filter the liquor through paper, which is to be kept in vessels closely stopped. The lime-water from calcined oyster-shells is prepared in the same manner.

Lime-water is principally used for the gravel; in which case, from a pint or two or more of it may be drank daily. Externally it is used for washing foul ulcers, and removing the itch, and other diseases of the skin.

Compound Lime Water.—Take shavings of guaiacum wood, half a pound; liquorice-root, one ounce; sassafras bark, half an ounce; coriander-seeds, three drachms; simple lime-water, six pints. Infuse without heat for two days, and then strain off the liquor.

In the same manner may lime-water be impregnated with the virtues of the other vegetable substances. Such impregnation not only renders the water more agreeable to the palate, but also a more efficacious medicine, especially in cutaneous disorders and foulness of the blood and juices.

It may be taken in the same quantity as the simple water.

Sublimate Water.—Dissolve eight grains of the corrosive sublimate in a pint of cinnamon-water. If a stronger solution be wanted, a double or triple quantity of sublimate may be used.

The principal intention of this is to cleanse foul ulcers, and consume proud flesh.

Styptic Water.—Take of blue vitriol and alum, each an ounce and a half; water, one pint. Boil them until the salts are dissolved, then filter the liquor, and add to it a drachm of the oil of vitriol.

This water is used for stopping a bleeding at the nose, and other hæmorrhages; for which purpose cloths or dossils dipped in it must be applied to the part.

Tar Water.—Pour a gallon of water on two pounds of Norway tar, and stir them strongly together with a wooden rod; after they have stood to settle for two days, pour off the water for use.

Though tar-water falls greatly short of the character which has been given of it, yet it possesses some medicinal virtues. It sensibly raises the pulse, increases the secretions, and sometimes opens the body, or occasions vomiting.

A pint of it may be drank daily, or more, if the stomach can bear it. It is generally ordered to be taken on an empty stomach, viz. four ounces morning and evening, and the same quantity about two hours after breakfast and dinner.

SIMPLE DISTILLED WATERS.

A GREAT number of distilled waters were formerly kept in the shops, and are still retained in some Dispensatories. But we consider them chiefly in the light of grateful diluents, suitable vehicles for medicines of greater efficacy, or for rendering disgusting

* See *Collyrium of Dead*.

diluents, suitable vehicles for medicines of greater efficacy, or for rendering disgusting ones more agreeable to the palate and stomach. We shall therefore insert only a few of those which are best adapted to these intentions.

The management of a still being now generally understood, it is needless to spend time in giving directions for that purpose.

Cinnamon Water.—Steep one pound of cinnamon bark, bruised, in a gallon and a half of water, and one pint of brandy, for two days; and then distil off one gallon.

This is an agreeable aromatic water, possessing in a high degree the fragrance and cordial virtues of the spice.

Pennyroyal Water.—Take of penny-royal leaves, dried, a pound and a half; water, from a gallon and a half to two gallons. Draw off by distillation one gallon.

This water possesses, in a considerable degree, the smell, taste, and virtues of the plant. It is given in mixtures and juleps to hysteric patients. An infusion of the herb in boiling water answers nearly the same purposes.

Peppermint Water.—This is made in the same manner as the preceding.

Spearmint Water.—This may also be prepared in the same way as the penny-royal water.

This and the preceding are useful stomachic waters, and will sometimes relieve vomiting, especially when it proceeds from indigestion, or cold viscid phlegm. They are likewise useful in some colicky complaints, the gout in the stomach, &c., particularly the peppermint-water.—An infusion of the fresh plant is frequently found to have the same effects as the distilled water.

Rose Water.—Take of roses, fresh gathered, six pounds; water, two gallons. Distil off one gallon.

This water is principally valued on account of its fine flavour.

Jamaica-pepper Water.—Take of Jamaica pepper, half a pound; water, a gallon and a half. Distil off one gallon.

This is a very elegant distilled water, and may in most cases supply the place of the more costly spice waters.

SPIRITUOUS DISTILLED WATERS.

Spirituos Cinnamon Water.—Take of cinnamon bark, one pound; proof spirit, and common water, of each one gallon. Steep the cinnamon in the liquor for two days; then distil off one gallon.

Spirituos Jamaica-pepper Water.—Take of Jamaica pepper, half a pound; proof spirit, three gallons; water, two gallons. Distil off three gallons.

This is a sufficiently agreeable cordial, and may supply the place of the *Aromatic Water*.

WHEYS.

Alum Whey.—Boil two drachms of powdered alum in a pint of milk, till it is curdled; then strain out the whey.

This whey is beneficial in an immoderate flow of the *menses*, and in a *diabetes*, or excessive discharge of urine.

The dose is two, three, or four ounces, according as the stomach will bear it, three times a-day. If it should occasion vomiting, it may be diluted.

Mustard Whey.—Take milk and water, of each a pint; bruised mustard-seed, an ounce and a half. Boil them together till the curd is perfectly separated; afterwards strain the whey through a cloth.

This is the most elegant, and by no means the least efficacious method of exhibiting mustard. It warms and invigorates the habit, and promotes the different secretions. Hence, in the low state of *nervous fevers*, it will often supply the place of wine. It is also of use in the chronic rheumatism, palsy, dropsy, &c. The addition of a little sugar will render it more agreeable. The dose is an ordinary tea-cupful four or five times a-day.

Scorbutic Whey.—This whey is made by boiling half a pint of the scorbutic juices in a quart of cow's milk. More benefit, however, is to be expected from eating the plants, than from their expressed juices.

The scorbutic plants are, bitter oranges, brooklime, garden scurvy-grass, and water-cresses. A number of other wheys may be prepared nearly in the same manner, as orange-whey, cream-of-tartar-whey, &c. These are cooling pleasant drinks in fevers, and may be rendered cordial, when necessary, by the addition of wine.

WINES.

The effects of wine are, to raise the pulse, promote perspiration, warm the habit, and exhilarate the spirits. The red wines, besides these effects, have an astringent quality, by which they strengthen the tone of the stomach and intestines, and by this means prove serviceable in restraining immoderate secretions. The thin sharp wines have a different tendency. They pass off freely by the different excretories, and gently open the body. The effects of the full-bodied wines are, however, much more durable than those of the thinner. All sweet wines contain a glutinous substance, and do not pass off freely. Hence they will heat the body more than an equal quantity of any other wine, though it should contain fully as much spirit. From the obvious qualities of wine, it must appear to be an excellent cordial medicine. Indeed, to say the truth, it is worth all the rest put together. But to answer this character, it must be sound and good. No benefit is to be expected from the common trash that is often sold by the name of wine, without possessing one drop of the juice of the grape. Perhaps no medicine is more rarely obtained genuine than wine. Wine is not only used as a medicine, but is also employed as a *menstruum* for extracting the virtues of other medicinal substances; for which it is not ill adapted, being a compound of water, inflammable spirit and acid; by which means it is enabled to act upon vegetable and animal substances, and also to dissolve some bodies of the metallic kind, so as to impregnate itself with their virtue, as steel, antimony, &c.

Anthelmintic Wine.—Take of rhubarb, half an ounce; worm-seed, an ounce. Bruise them, and infuse without heat in two pints of red Port wine for a few days, then strain off the wine.

As the stomachs of persons afflicted with worms are always debilitated, red wine alone will often prove serviceable. It must, however, have still better effects when joined with bitter and purgative ingredients, as in the above form.—A glass of this wine may be taken twice or thrice a-day.

Antimonial Wine.—Take glass of antimony, reduced to a fine powder, half an ounce; Lisbon wine, eight ounces. Digest, without heat, for three or four days, now and then shaking the bottle; afterwards filter the wine through paper.

The dose of this wine varies according to the intention. As an alterative and diaphoretic, it may be taken from ten to fifty or sixty drops. In a large dose it generally proves cathartic, or excites vomiting.

Bitter Wine.—Take of gentian-root, yellow rind of lemon-peel, fresh, each one ounce; long pepper, two drachms; mountain wine, two pints. Infuse without heat for a week, and strain out the wine for use.

In complaints arising from weakness of the stomach, or indigestion, a glass of this wine may be taken an hour before dinner and supper.

Ipecacuanha Wine.—Take of ipecacuanha, in powder, one ounce, mountain wine, a pint. Infuse for three or four days; then filter the tincture.

This is a safe vomit, and answers extremely well for such persons as cannot swallow the powder, or whose stomachs are too irritable to bear it. The dose is from one ounce to an ounce and a half.

Chalybeate, or Steel Wine.—Take filings of iron, two ounces; cinnamon and mace, of each two drachms; Rhenish wine, two pints. Infuse for three or four weeks, frequently shaking the bottle; then pass the wine through a filter.

In obstructions of the *menses*, this preparation of iron may be taken, in the dose of half a wine-glass twice or thrice a-day.

The medicine would probably be as good if made with Lisbon wine, sharpened with half an ounce of the cream of tartar, or a small quantity of the vitriolic acid.

Stomach Wine.—Take of Peruvian bark, grossly powdered, an ounce; cardamom-seeds, and orange-peel, bruised, of each two drachms. Infuse in a bottle of white Port or Lisbon wine for five or six days; then strain off the wine.

This wine is not only of service in debility of the stomach and intestines, but may also be taken as a preventive, by persons liable to the intermittent fever, or who reside in places where this disease prevails. It will be of use likewise to those who recover slowly after fevers of any kind, as it assists digestion, and helps to restore the tone and vigour of the system.

A glass of it may be taken two or three times a-day.

Preparation of the Lichen Islandicus.

THE lichen islandicus has of late been much extolled as a remedy in pulmonary consumption. Not that it is supposed to possess any peculiar virtues as a pectoral medi-

cine, but is rather to be considered as a nutritious vegetable jelly, which promotes digestion by its slight bitterness, and at the same time sufficiently supports the strength without stimulating the system; thus affording the powers of the constitution an opportunity of subduing the diseased action of the lungs. Many animals become very fat while living solely on the lichen islandicus, a proof that it affords abundant nutriment. To derive any essential benefit from the use of this article, it must constitute the sole subsistence. Persons who possess sufficient resolution to persist in such a regimen will find the lichen, prepared according to the following directions, an agreeable article of food:—

Four ounces of the lichen, cut in small pieces, are to be boiled in three pints of water, till the quantity be reduced to two. To this add a tea-spoonful of fine wheaten flour, and four ounces of refined sugar; take sixty sweet almonds, and twenty-four bitter ones, blanched; beat them into a paste, with a small quantity of water, mingle this with the decoction, and then strain through a linen cloth, previously moistened with warm water. By this process, a nutritious vegetable jelly is formed, which may be poured into moulds, or glasses, and eaten alone, or with milk. The aromatic flavour of the almonds blends itself with, and completely conceals the peculiar bitter of the lichen, and forms a very palatable, easily-digested, and nutritive aliment. A. P. B.

CONTENTS.

CHAP. I.		The Kind of Exercise recommended to Studious Habits	
OBSERVATIONS ON DIET, particularly that of the common people - -	Page 19		54
General Observations on Aliment - - -	20	CHAP. III.	
Of Bread - - -	21	NON-NATURALS - -	55
Boiled Grain - - -	24	Observations on Diet -	id.
Butter - - -	27	Vegetable Diet - -	57
Fruits and Roots - -	29	Plain Rules in the Selection of Diet, &c. - -	59
Broths and Soups - -	34	General Observations on Diet, long Fasting, its consequences, Regularities, &c.	60
Remarks on Food - -	37	Suppers and Breakfasts contrasted, &c. - -	61
Food considered in a medical point of view - -	39	CHAP. IV.	
Differences between Vegetable and Animal Food -	id.	AIR - - - -	62
CHAP. II.		Burying within Churches, a reprehensible Practice	63
ON SEDENTARY, STUDIOUS, and LABORIOUS OCCUPATIONS, &c.	40	Stagnated Air - -	64
On various Employments	42	Air of great Towns -	65
Soldiers and Sailors, the consequences of their Employment - - -	44	CHAP. V.	
The Sedentary - - -	46	EXERCISE - - - -	66
Injurious Effect of long Inclined Posture, &c. -	47	Inactivity, its consequences	67
Advice to the Sedentary, &c.	49	Exercise in the open Air	68
The Sedentary Studious	50	Indolence, its consequences	69
Diseases to which Studious People are more peculiarly liable - - -	id.	CHAP. VI.	
Advice to Studious and Intense Thinkers - - -	52	SLEEP, its benefits, &c. -	69
Desk Occupations, &c.	53	To procure refreshing Sleep, &c. - - -	70
Morning best adapted for Study and Exercise, &c. -	id.	Early rising - - -	71
Effects of Music on the Mind	id.	CHAP. VII.	
Dr. Buchan's Plan of invigorating the Spirits -	54	CLOTHING, to be suited to the climate and season of the year	71
Diet of the Studious -	id.	CHAP. VIII.	
		INTEMPERANCE, how hurtful	75
		CHAP. IX.	
		CLEANLINESS - - -	78
		Necessity of frequent Ablutions - - -	80

CHAP. X.		CHAP. V.	
INFECTION and CONTAGION, what, and how communicated	82	CONTINUED FEVERS	120
CHAP. XI.		Synochus or Simple Continued Fever. Symptoms, Medical Treatment, Diet, &c.	120—122
AFFECTIONS of the MIND	85	CHAP. VI.	
Anger	86	INFLAMMATORY, ACUTE or AR- DENT FEVER (<i>Synocha</i>)	122
Fear	id.	Causes, Symptoms, Treatment, Diet, &c.	122—127
Effects of Fear on <i>Child-bed</i> <i>Women</i>	id.	CHAP. VII.	
Reprehensible Customs	87	SLOW or NERVOUS FEVER (<i>Ty- phus Mitior</i>)	127
Grief	88	Causes, Symptoms, Diet, Med- ical Treatment, &c.	127—133
Love	90	CHAP. VIII.	
Religious Melancholy	id.	MALIGNANT and PUTRID, or SPOT- TED FEVER (<i>Typhus gravior</i>)	133
CHAP. XII.		Causes, Symptoms, Diet, Med- ical Treatment, &c.	133—140
THE NATURAL EVACUATIONS	91	CHAP. IX.	
The Faecal Evacuations, &c.	id.	MILIARY FEVER. Causes, Symp- toms, Regimen, Medicine, &c.	140—143
Urine	92	CHAP. X.	
Perspiration	94	BILIARY FEVER, &c.	143
Causes of obstructed Perspi- ration	id.	CHAP. XI.	
Wet Clothes	95	PLEURISY (<i>Pleuritis</i>). Causes, Symptoms, Diet, Medicine, &c.	144—148
Wet Feet	id.	Bastard Pleurisy, &c.	149
Night Air	id.	Paraphrenitis, or Inflammation of the Diaphragm	id.
Damp Beds	id.	CHAP. XII.	
Damp Houses	96	PERIPNEUMONY, or Inflammation of the Lungs, (<i>Pneumonia</i>), Causes, Symptoms, Diet, &c.	150—151
Sudden Transitions, &c.	97	CHAP. XIII.	
PART II.		PHTHISIS, or Pulmonary Con- sumption. (<i>Phthisis Pulmo- nalis</i>). Causes, Symptoms, Diet, Medicine, &c.	152—160
CHAP. I.		Nervous Consumption	160
THE KNOWLEDGE and CURE of DISEASES	100	Prevention of Consumption or Atrophy	id.
Definition of Diseases, &c.	id.		
CHAP. II.			
GENERAL OBSERVATIONS on FE- VERS	104		
Causes of Fevers	id.		
Fever considered as an Effort of Nature, &c.	105		
Symptoms of Fever	106		
Treatment of Fevers, &c.	108		
CHAP. III.			
INTERMITTENT FEVERS or AGUES, Causes, Symptoms, Stages, Medical Treatment, Diet, &c.	109		
CHAP. IV.			
REMITTENT FEVER, Cause, Symp- toms, Diet, Medical Treat- ment	117		

On the Means of Preventing Pulmonary Consumption 161	Inflammation of the Kidneys, &c. - - - 217
CHAP. XIV.	Inflammation of the Bladder, &c. - - - 219
SMALL POX. (<i>Variola</i> .) Causes, Symptoms, Diet, Medicine 168--176	Inflammation of the Liver, &c. 220
CHAP. XV.	CHAP. XXIII.
MEASLES. (<i>Morbilli</i> or <i>Rubeola</i> .) Causes, Symptoms, Regimen, Medical Treatment, &c. 176--179	CHOLERA MORBUS, and other ex- cessive Discharges from the Stomach and Bowels. Causes, Symptoms, Medicine, &c. - - - 223--226
CHAP. XVI.	Diarrhœa, or Looseness 226
SCARLET FEVER. (<i>Scarlatina</i>) 179	Vomiting, from various causes 228
CHAP. XVII.	CHAP. XXIV.
ERESIPELAS, or ST. ANTHONY'S FIRE. Causes, Symptoms, Treatment, Diet, &c. 181-- 185	DIABETES, and other Disorders of the Urinary Organs. Causes, Symptoms, Regi- men, Medicine 230--232
CHAP. XVIII.	Suppression of Urine 233
INFLAMMATION of the BRAIN. (<i>Phrenitis</i> .) - 185--189	Gravel and Stone, &c. &c. 234
CHAP. XIX.	CHAP. XXV.
OPHTHALMIA, or Inflammation of the Eyes. Causes, Symptoms, Regimen, Medical Treatment 189--193	INVOLUNTARY DISCHARGES OF BLOOD - - - 238
CHAP. XX.	Bleeding at the Nose 239
QUINSEY, or Inflammatory Sore Throat, (<i>Cynanche Tonsilla- ris</i> .) Causes, Symptoms, Treatment, &c. 193--197	Bleeding and Blind Piles 241
Malignant Quinsey, or Putrid Ulcerous Sore Throat, (<i>Cy- nanche Maligna</i> .) Causes, Symptoms, Regimen, Medi- cine, &c. - 197--201	Spitting of Blood - 244
Mumps, (<i>Cynanche Parotidæa</i>) 201	Vomiting of Blood - 247
CHAP. XXI.	Bloody Urine - - 248
COLDS and COUGHS, (<i>Catarrhal Affections</i>) - - 202	Dysentery, or Bloody Flux 249
Common Cough - 205	CHAP. XXVI.
Hooping Cough, or Chin Cough (<i>Pertussis</i>) - 207--209	HEAD-ACH - - - 254
CHAP. XXII.	Causes, Symptoms, &c. id.
INFLAMMATION of the STOMACH, (<i>Gastritis</i> .) Causes, Sym- ptoms, Regimen, Medicine 210	Tooth-ach - - - 156
Inflammation of the Intestines, &c. - - - 211	Ear-ach - - - 258
	Pain of the Stomach, &c. 259
	CHAP. XXVII.
	WORMS. Causes, Symptoms, &c. 261
	CHAP. XXVIII.
	JAUNDICE. Causes, Symptoms, Regimen, Medicine, &c. 265
	CHAP. XXIX.
	DROPSY. Causes, Symptoms, Di- et, Medical Treatment, Varie- ties - - - 268
	CHAP. XXX.
	GOUT. Causes, Symptoms, Re- gimen, Medicine, &c. 273
	Rheumatism, &c. - 277

CHAP. XXXI.			
SCURVY. Causes, Symptoms,			
Cure, &c.	-	281	
Scrofula, or King's Evil	-	285	
	-	290	
Itch	-	290	
Asthma	-	292	
Apoplexy	-	296	
Costiveness and other Affec-			
tions of the Stomach and			
Bowels	-	299	
Want of Appetite	-	300	
Heartburn	-	301	
Nervous Diseases	-	303	
Melancholy	-	306	
Palsy	-	309	
Epilepsy, or Falling Sickness			
	-	311	
St. Vitus's Dance	-	313	
Hiccup	-	315	
Cramp of the Stomach	-	316	
Sardonic Laugh	-	317	
Nightmare	-	id.	
Swooning	-	318	
Flatulencies	-	320	
Low Spirits	-	321	
Hysteric Affections	-	322	
Hypochondriac Affections	-	325	
CHAP. XXXII.			
DISORDER of the SENSES	-	329	
Of the Eye	-	id.	
Gutta Serena, or Amaurosis	-	330	
Cataract	-	id.	
Myopia, or Shortsightedness	-	331	
Squinting	-	id.	
Spots or Specks of the Eyes	id.		
Blood-shot Eye	-	id.	
Watery or Weeping Eye	id.		
The Ear	-	332	
Taste and Smell	-	333	
Touch	-	335	
CHAP. XXXIII.			
SCIRRHRUS and CANCER. Causes,			
Symptoms, Treatment, &c.	-	336	
CHAP. XXXIV.			
POISONS.	-	342	
Mineral Poisons. Symptoms,			
Treatment, Tests, &c.	-	342	
	-	348	
Hydrophobia	-	348	
Poisonous Animals and Plants	-	350	
Viper, Rattlesnake, &c.	id.		
PART III.			
CHAP. I.			
SURGERY	-	352	
Bleeding	-	id.	
Topical Blood-letting	-	355	
Inflammations and Abscesses	-	id.	
Wounds	-	357	
Burns	-	359	
Mr. Cleghorn's plan of treat-			
ing Burns and Scalds	-	361	
Sir James Earle's Plan	id.		
Bruises	-	362	
ULCERS	-	id.	
Cures of Ulcers by Roller and			
Compresses, &c. on Mr.			
Whately's Plan	-	363	
Mr. Baynton's plan of curing			
old Ulcers of the Leg by			
means of adhesive Plaster,			
without rest	-	365	
Fistula in Ano	-	367	
CHAP. II.			
DISLOCATIONS	-	369	
Dislocations of the Jaw	-	370	
Neck	-	id.	
Ribs	-	371	
Shoulder	-	id.	
Elbow	-	372	
Clavicle, or Collar Bone	id.		
Patella, or Knee Pan	-	373	
Thigh	-	id.	
CHAP. III.			
BROKEN BONES	-	374	
CHAP. IV.			
OF STRAINS	-	377	
Ruptures	-	id.	
CHAP. V.			
CASUALTIES	-	378	
Substances stopped between			
the Mouth and Throat	-	379	
Suspended Animation and Re-			
suscitation	-	382	
Noxious Vapours, or Aerial			
Poisons	-	385	

CONTENTS.

xxxvii

Effects of Extreme Cold	387	Stages of Labour	-	412
Heat	388	Symptoms preceding Labour		
Fainting Fits	-	id.		
Intoxication	-	accompanying La-		
Of Suffocation and Strangulation,		bour	-	413
from Hanging, Drowning, and other Causes	393	Process of Natural Labours		414
Persons who expire in Convulsion Fits	395	Management of the After-bur-		
		den	-	415
		Tedious Labour	-	416
		Treatment after Delivery	id.	
		After-pains	-	417
		Costiveness	-	id.
		Inflammation of the Womb	id.	
		Suppression of the Lochia		418
		Inflammation of the Breast	id.	
		Fretted or chapped Nipples	id.	
		Miliary Fever	-	id.
		Puerperal Fever	-	419
		Milk Fever	-	420
CHAP. VI.		CHAP. XI.		
DISEASES OF WOMEN	397	BARRENNESS	-	421
The Menstrual Discharge	id.			
Immoderate Flow of the Menses	-			
Fluor Albus	-	CHAP. XII.		
Difficult Menstruation	403	DISEASES OF INFANTS	-	422
Cessation of the Menses (commonly called the turn of life)	id.	Meconium	-	423
		Aphthæ and Thrush		id.
		Hiccups	-	424
		Acidities	-	id.
		Galling and Excoriation		425
		Stoppage of the Nose		426
		Vomiting	-	id.
		Looseness	-	427
		Cutaneous Eruptions		428
		Of Croup	-	429
		Teething	-	431
		Rickets	-	433
		Convulsions	-	435
		Water in the Head	-	436
		CHAP. XI.		
		OBSERVATIONS ON THE MORAL AND		
		PHYSICAL MANAGEMENT OF		
		CHILDREN, DISEASES, &c.		438
		Of Diseased Parents		441
		Clothing of Children		442
		Food of Children	-	446
		Exercise of Children		449
		Bad Effects of unwholesome		
		Air upon Children		454
		Nurses	-	456
		CHAP. XII.		
		COLD BATHING in fresh and salt		
CHAP. VII.				
DISEASES OF PREGNANCY	403			
Nausea and Vomiting	404			
Heartburn	-			
Head-ach with Plethora	id.			
Costiveness and Piles	id.			
Tooth-ach	-			
Longings	-			
Hysterics	-			
Diarrhœa	-			
Suppression of Urine	id.			
Troublesome Itchings	id.			
Puffy Swellings of the Feet	id.			
Cramps of the Legs & Thighs	407			
Restlessness & Want of Sleep	id.			
Varicose Veins	-			
Jaundice	-			
Incontinency of Urine	id.			
Over-distention of the Skin	408			
False Pains	-			
CHAP. VIII.				
CONVULSIONS	-			
CHAP. IX.				
ABORTION	-			
CHAP. X.				
MANAGEMENT OF CHILD-BED Wo-				
MEN	-			
Parturition	-			
Classification of Labours	id.			

water, with remarks on the
cases in which the Warm
Bath is more advisable 459
Effects of Cold Bathing, &c. 463

Cautions to be observed 464

CHAP. XIII.

MINERAL WATERS - 468

Classification of Mineral Wa-
ters - 470

Tunbridge Wells - 471

Pymont and German Spa
473

Cheltenham Waters - 475

Scarborough - 476

Seltzer - 477

Harrowgate - 478

Moffat - 481

Bath Hot Springs - id.

Buxton - 484

Matlock - 485

Bristol Hot Well - id.

Continental Hot Springs 487

Malvern Water - 488

Aluminous Chalybeate Spring
in the Isle of Wight 489

CHAP. XIV.

Vegetable Poisons - 490

Narcotic Poisons - 491

Poisonous Mushrooms 492

CHAP. XV.

POISONOUS FISH - 492

CHAP. XVI.

VACCINATION - 493

CHAP. XVII.

SEA-SICKNESS - 494

Imperial Marine Tincture
for - 495

APPENDIX.

MEDICINES USED IN PRACTICE,
&c. - - - iii

DOMESTIC PHARMACOPEIA - ix

Balsams - - - id.

Boluses - - - id.

Cataplasms - - - x

Clysters - - - id.

Collyria, or Eye Water - xi

Confections - - - xii

Conserves and Preserves id.

Decoctions - - - id.

Draughts - - - xiii

Electuaries - - - xiv

Emulsions - - - xv

Extracts - - - id.

Fomentations - - - xvi

Gargles - - - id.

Infusions - - - xvii

Juleps - - - xviii

Mixtures - - - id.

Ointments - - - xix

Liniments - - - id.

Cerates - - - id.

Pills - - - xxi

Plasters - - - xxii

Poultices - - - xxiv

Powders - - - id.

Syrups - - - xxv

Tinctures - - - xxvi

Elixirs - - - id.

Vinegars - - - xxviii

Waters by Infusion - xxix

Simple Distilled Waters - id.

Spirituos Distil'd Waters xxx

Wheys - - - id.

Wines - - - xxxi

Preparation of the Lichen

Islandicus - - - id.

DFG

Blood, involuntary discharges of - - -	238	C.	
—, vomiting of - - -	247	Cancer - - -	336
—, spitting of - - -	244	Candied orange peel - -	xii
Boluses - - -	ix	—, lemon ditto - - -	<i>id.</i>
Bolus, astringent - -	<i>id.</i>	Carrots to be eaten young	33
—, diaphoretic - - -	<i>id.</i>	Casualties - - -	378
—, mercurial - - -	x	Catamenia. <i>See</i> Discharge, Menstrual.	
—, of rhubarb and mercury - - -	<i>id.</i>	Cataplasms - - -	x
—, pectoral - - -	<i>id.</i>	—, discutient, - - -	<i>id.</i>
—, purging - - -	<i>id.</i>	—, ripening - - -	<i>id.</i>
Bones, broken - - -	374	Cataract - - -	330
—, treatment of - - -	<i>id.</i>	Celsus, his justly celebrated advice - - -	99
—, setting of - - -	375	Cheltenham waters - - -	475
Bowels, excessive discharges from. <i>See</i> Cholera Morbus.		Child-bed women, management of - - -	411
Brain, dropsy of - - -	436	Children, moral and physical management of - -	438
—, inflammation of - -	185	—, diseases of - - -	<i>id.</i>
Bread, considered as food	21	—, mothers, advice to, concerning - -	439
—, often spoiled to please the eye - - -	22	—, seldom well unless their bodies are kept gently open - - -	26
—, made of different kinds of grain - - -	23	Chimney sweeper's cancer - - -	78
—, the greatest consumption of, occasioned by tea - - -	<i>id.</i>	—, Pott, Mr. quoted on - -	<i>id.</i>
Breakfasts and suppers contrasted - - -	61	—, Earl, Mr. quoted on - -	79
Breast, inflammation of -	418	Chin-cough. <i>See</i> Cough, Hooping.	
Bristol hot-wells - - -	485	Choking - - -	379
Broths and soups - - -	34	Choke-damp explained - -	41
—, a dish of great antiquity - - -	36	Cholera morbus - - -	223
—, Hufiland, Dr. quoted on soups - - -	37	Clarke, Sir Arthur, hypothesis, faulty, relative to immersion in cold water	98-99
Bruises - - -	362	Cleanliness - - -	78
Buchan, Dr. his plan of exercise - - -	54	Clothing, how to be regulated - - -	71
Buckwheat, highly nutritious	27	—, to be suited to climate - - -	<i>id.</i>
Burns - - -	359	—, to the season of the year - - -	72
—, Cleghorn, Mr. his plan of treating - - -	361	—, how hurtful - - -	<i>id.</i>
—, Kentish, Mr. his plan - -	360	—, tight bandages to be avoided in - - -	73
—, Earle, Sir James, his plan - - -	361	—, of children - - -	442
Burying within churches, the practice condemned - -	63	—, necessary regulations and observations on - -	445
—, its practice among the Jews, Romans, &c. -	<i>id.</i>	Clysters - - -	x
Butter, considered as food	27	Clyster, emollient - - -	xi
Buxton waters - - -	484	—, laxative - - -	<i>id.</i>

Clyster, carminative	xi	Deadly nightshade. <i>See</i> Poisons Narcotic.	
—, oily	<i>id.</i>	Decoctions	xii
—, starch	<i>id.</i>	— of mallows	xiii
—, turpentine	<i>id.</i>	— common	<i>id.</i>
—, vinegar	<i>id.</i>	— logwood	<i>id.</i>
Cold, the effects of extreme	387	— bark	<i>id.</i>
Colds, more destructive than plagues	72	— compound of	<i>id.</i>
— Boerhaave, quoted on	74	— sarsaparilla	<i>id.</i>
Colds and coughs	202	— seneka	<i>id.</i>
Colic	214	— white	<i>id.</i>
Collyria	xi	Delivery, treatment after	416
Collyrium of alum	<i>id.</i>	Dentition. <i>See</i> Teething.	
— vitriolic	<i>id.</i>	Desk occupations	53
— of lead	<i>id.</i>	Diabetes	230
Confection, japonic	xii	— Doctors Farrier and Rollo, quoted on	232
— of catechu	<i>id.</i>	Diet, observations on	55
Conserves	<i>id.</i>	— vegetable and animal	57
Conserve of red roses	<i>id.</i>	—, observations on, &c.	19
— of sloe	<i>id.</i>	—, plain rules for the selection of	59
Consumption, pulmonary	152	—, general observations on	60
— on the means		—, sudden changes in	61
— of preventing	161	Diarrhœa	226
Convulsions, during pregnancy or labour	408	— of pregnant women	406
Convulsion fits, persons who expire in	395	Discharge, menstrual	401
Convulsions of infants	435	Diseases of the extremities, how often caused	43
Copper and its preparations. <i>See</i> Mineral Poisons.		—, knowledge and cure	100
Corn, Indian, when boiled	26	—, females liable to some which do not affect the opposite sex	101
Corrosive sublimate. <i>See</i> Mineral Poisons.		—, particular constitutions disposed to	<i>id.</i>
Costiveness	299	Dislocations	369
—, after delivery	417	— of the jaw	370
Cough, hooping	207	— neck	<i>id.</i>
Cramp, of the stomach	316	— ribs	371
Cramps of the legs & thighs of pregnant women	407	— shoulder	372
Croup	429	— elbow	<i>id.</i>
— Turnbull, Dr. William, quoted on	430	— clavicle or collar bone	<i>id.</i>
Customs, reprehensible ones	87	— patella or kneepan	373
		— thigh	<i>id.</i>
D.		Draughts	xiii
Daffey's Elixir, a dangerous nostrum in the nursery	457	Draught, anodyne	xiv
Dalby's carminative, a dangerous nostrum in the nursery	<i>id.</i>	— diuretic	<i>id.</i>
Damp beds	95	— purging	<i>id.</i>
— houses	96	— sweating	<i>id.</i>
		— vomiting	<i>id.</i>
		Dress, the perfection of, in	

what consists - - -	73	Exercise, as necessary as food - - -	66
Dress, military smartness in, or adorned - - -	74	—, to be taken in the open air - - -	68
Drinking, how health injured by it - - -	77	—, glandular obstructions removed by - - -	66
—, to excess, the barbarous custom of - - -	<i>id.</i>	Exercise of children - - -	449
—, hard, increase of consumption imputed to - - -	<i>id.</i>	—, the kind of, recommended for studious habits - - -	54
Dropsy - - -	268	Extracts - - -	xv
— of the extremities, or anasarca - - -	272	Evacuations, the natural - - -	91
— of the abdomen, or ascites - - -	<i>id.</i>	Eye, disorders of - - -	329
Drowned persons - - -	382	—, spots or specks on - - -	331
Drugs necessary for private practice - - -	iii	—, blood-shot - - -	<i>id.</i>
Dysentery - - -	249	—, watery or weeping - - -	<i>id.</i>
— putrid - - -	251	Eye-waters. <i>See</i> Collyria.	
		Eyes, inflammation of - - -	189
		F.	
E.		Falling sickness - - -	311
Ear-ach - - -	258	Fasting, long, its consequences - - -	60
Ear, disorder of - - -	332	Fear - - -	86
Electuaries - - -	xiv	—, its effects on child-bed women - - -	<i>id.</i>
Electuary, lenitive - - -	<i>id.</i>	Feet, they suffer by pressure - - -	72
— for dysentery - - -	<i>id.</i>	Fever, intermittent - - -	109
— for epilepsy - - -	<i>id.</i>	—, remittent, &c. - - -	117
— for obstructed menses - - -	xv	—, putrid remitting, of Bengal, Dr. Lind quoted on - - -	118
— of bark - - -	<i>id.</i>	—, continued - - -	120
— for piles - - -	<i>id.</i>	—, inflammatory, acute, or ardent - - -	122
— palsy - - -	<i>id.</i>	—, slow or nervous - - -	127
— rheumatism - - -	<i>id.</i>	—, malignant and putrid, or spotted fevers - - -	133
Elixir, paregoric - - -	xxvii	—, miliary - - -	140, 418
—, sacred - - -	<i>id.</i>	—, bilious - - -	143
—, stomachic - - -	<i>id.</i>	—, scarlet - - -	179
—, acid of vitriol - - -	xxviii	—, puerperal - - -	418
Employments, various observations on - - -	42	—, milk - - -	420
Emulsions - - -	xv	Fevers, general observations on - - -	104
—, common - - -	<i>id.</i>	—, beverage, in inflammatory - - -	106
—, arabic - - -	<i>id.</i>	—, exciting causes of, &c. - - -	109
—, camphorated - - -	<i>id.</i>	—, treatment, &c. - - -	110
—, of gum ammoniac - - -	<i>id.</i>	—, Hoffman and Cul-len on - - -	106
—, oily - - -	<i>id.</i>	—, hypothesis of Stahl - - -	<i>id.</i>
Epilepsy. <i>See</i> Falling Sickness.			
Erysipelas, or St. Anthony's fire - - -	181		
Eruptions, cutaneous of children - - -	428		
Excoriation. <i>See</i> Galling.			

INDEX.

xliii

Fewillea cordifolia, the fruit of, an antidote, &c.	491
Fish, poisonous	492
Fistula in ano	367
——, lachrymalis	331
Flatulencies	320
Fits, fainting	389
Fluor albus	401
Flux, bloody. <i>See</i> Dysentery.	
Fomentations	xvi
——, anodyne	<i>id.</i>
——, aromatic	<i>id.</i>
——, emollient	<i>id.</i>
——, strengthening	<i>id.</i>
Food of children	446
——, how to be regulated	448
—— considered in a medical point of view	39
Foxglove. <i>See</i> Poison Narcotic.	
Fruits and roots, substitutes for bread	29
Fumigations, disinfecting	84

G.

Galling	425
Gargles	xvi
Gargle, attenuating	<i>id.</i>
——, common	<i>id.</i>
——, detergent	<i>id.</i>
——, emollient	xvii
Gluttony, &c. ought to be checked	78
Godfrey's cordial, a dangerous nostrum in the nursery	457
Gout	273
Grain, boiled	24
Gravel and stone	234
Grief, the most destructive of all passions	88
Ground, cultivation of, conducive to health	49
Gutta serena	330

H.

Hanging. <i>See</i> Suffocation.	
Harrowgate water	478
Head-ach, with plethora of pregnant women	405
Head-ach	254

Heart-burn	301
Heart-burn of pregnant women	405
Heat, the effects of extreme	388
Hernia. <i>See</i> Ruptures.	
Henbane. <i>See</i> Poisons narcotic.	
Hiccup	315
Hiccups, of infants	424
Hiera picra. <i>See</i> Tincture sacred and sacred elixir.	
Hossack, Dr. of New York, quoted	82
Hot-springs, continental	487
Humane Society, Royal	396
Hydrophobia	348
Hypochondriac affections	325
Hypochondriacs, Cullen's description of	52
Hypochondriasis. <i>See</i> Low Spirits.	
Hysteria. <i>See</i> Hysteric affections.	
Hysteric affections	322
Hysterics of pregnant women	406

I.

Ideas, a change of, necessary to health	89
Inactive, the, always complaining, of what	67
Indolence, its consequences	69
Infants, diseases of	422
—— arise chiefly from the bowels	<i>id.</i>
Infection and contagion	82
—— observations and opinions on	<i>id.</i>
——, diseases, many caused by	<i>id.</i>
—— cautions against	84
Inflammations, &c.	355
Inflammation of the bladder. <i>See</i> Bladder.	
—— of the brain. <i>See</i> Brain.	
—— of the eyes. <i>See</i> Eyes.	
—— of the intestines. <i>See</i> Intestines.	
—— of the kidneys. <i>See</i> Kidneys.	

Inflammation of the liver.		King's evil	- - - 285
See Liver.			
----- of the lungs.		L.	
See Lungs.			
----- of the stom-		Laborious occupations, &c.	
ach. See Stomach.		remarks on	- - - 40
----- of the throat.		Labours, classification of	412
See Quinsey.		Labour, stages of	- - - <i>id.</i>
----- of the womb.		-----, symptoms preceding	<i>id.</i>
See Womb.		-----, symptoms accompa-	
Infusions - - - - xvii		nying	- - - 413
Infusion, bitter - - - <i>id.</i>		-----, process of, natural	414
----- of the bark - - - <i>id.</i>		-----, tedious - - -	416
----- Carduus Benedic-		-----, treatment after	<i>id.</i>
tus, or blessed thistle - <i>id.</i>		Laugh, sardonic - - -	317
----- linseed - - - <i>id.</i>		Lead and its preparations.	
----- of roses - - - <i>id.</i>		See Mineral Poisons.	
----- tamarinds and senna <i>id.</i>		Leeks - - - - -	34
----- Spanish - - - xviii		Lichen, islandicus, prepara-	
----- for the palsy - - - <i>id.</i>		tions of - - - - -	xxxii
----- antiscorbutic - - - <i>id.</i>		Lime. See Alkaline Earths.	
Intemperance - - - - 75		Liniment for burns - - -	xx
-----, the danger of	<i>id.</i>	-----, volatile - - -	<i>id.</i>
-----, a striking		-----, white - - -	<i>id.</i>
proof of afforded, by what	<i>id.</i>	-----, for piles - - -	<i>id.</i>
-----, families affect-		Liquors, fermented - - -	58
ed by it - - - - - 76		-----, observations on	59
Intestines, inflammation of		-----, hot, in warm rooms,	
the - - - - - 211		the danger of drinking	98
Intoxication, consequences		Liver, inflammation of	220
of - - - - - 76		Lochia, suppression of	418
----- hurtful to young		Longings of pregnant wo-	
persons - - - - - 78		men - - - - -	406
----- effects of, often		Looseness of infants - - -	427
fatal - - - - - 391		Love, perhaps the strongest	
Itch - - - - - 290		of all passions - - -	90
Itchings, troublesome, of		Low spirits - - - - - 321	
pregnant women - - - 406		Lungs, inflammation of	150
J.			
Jaundice - - - - - 265		M.	
----- of pregnant wo-		Malvern water - - - - 488	
men - - - - - 407		Marine tincture, imperial,	
Juleps - - - - - xviii		for sea-sickness and ner-	
Julep, camphorated - - - <i>id.</i>		vous disorders - - -	495
-----, cordial - - - - <i>id.</i>		Matlock waters - - -	485
-----, expectorating - - <i>id.</i>		Measles - - - - - 176	
-----, musk - - - - - <i>id.</i>		Mechanics, advice to - -	49
-----, saline - - - - - <i>id.</i>		Meconium - - - - - 423	
-----, vomiting - - - - <i>id.</i>		Medicines used in private	
K.		practice - - - - - iii-viii	
Kidneys, inflammation of	217	-----, names of - - - <i>id.</i>	
		-----, properties - - - <i>id.</i>	

INDEX.

xlv

Medicines, doses -	iii—viii	Oil, camphorated -	xxi
Melancholy -	306	Ointments -	xix
——, religious -	90	Ointment, yellow basilicum	id.
Menses, immoderate flow of	401	——, of calamine -	id.
——, cessation of -	403	——, emollient -	id.
Menstruation, difficult	id.	——, eye -	id.
Mercury and its preparations. See Mineral Poisons.		——, issue -	xx
Midwives, their ill-judged care of infants -	422	——, lead -	id.
Mind, the temper of in diseases, how to be attended to -	101	——, mercurial -	id.
——, affections of -	85	——, sulphur -	id.
Mindererus's spirit -	xxviii	——, for diseases of the skin -	id.
Mineral waters -	468	——, white -	id.
——, classifica-		Opium. See Poisons, Narcotic.	
tion of -	470	Onions how dressed -	34
Mixtures -	xviii	P.	
Mixture, astringent -	xix	Pain after -	417
——, diuretic -	id.	Pains false -	408
——, laxative, absorbent	id.	Palsy -	309
——, saline -	id.	Paraphrenitis -	149
——, squill -	id.	Parents, diseased, the effect upon children -	441
Moffat waters -	481	Parturition -	412
Mouth and nose, substances stopped between -	379	Peas and beans considered as food -	25
Mumps -	201	Perspiration. See Evacuations.	
Mushrooms, poisonous	492	——, causes of, obstructed -	94
Music, its effects on the mind -	53	Pills -	xxi
Myopia -	331	Pill, composing -	id.
N.		——, deobstruent -	id.
Nausea and vomiting of pregnant woman -	404	——, foetid -	id.
Nerves, weak, the companion of inactivity -	67	——, hemlock -	id.
Nervous diseases -	303	——, mercurial -	id.
Night-air -	95	——, mercurial sublimate	id.
Night-mare -	317	——, plumber's -	id.
Nipples, fretted or chapped	418	——, purging -	xxii
Nitre -	348	——, purgative -	id.
Non-naturals -	55	——, for the bile -	id.
Nose, bleeding at -	239	——, jaundice -	id.
——, stoppage of, in children -	426	——, stomachic -	id.
Nurses, the duty of -	456	——, squill -	id.
O.		——, strengthening -	id.
Oatmeal frequently made into bread -	25	Piles, blind and bleeding	241
		Placenta. See After Burden.	
		Plants, poisonous -	351
		Plasters -	xxii
		Plaster, common -	xxiii
		——, adhesive -	id.
		——, blistering -	id.

Plaster, gum - - -	<i>id.</i>	Rheumatism - - -	277
—, mercurial - - -	<i>id.</i>	Rickets - - -	433
—, stomach - - -	<i>id.</i>	Rising, early, a consequence	
—, warm - - -	<i>id.</i>	of what - - -	71
—, wax - - -	<i>id.</i>	Ruptures - - -	377
Pleurisy - - -	144		
—, bastard - - -	149	S.	
Poisons, mineral - - -	342	Sailors and soldiers, conse-	
—, vegetable - - -	342, 490	quences of their calling	44
—, animal - - -	342	Saint Anthony's fire, a dis-	
—, aërial. <i>See</i> Noxious		ease incident to labouring	
Vapours.		people - - -	43
—, narcotic - - -	491	—, iliac pas-	
Poor living after the course		sion, colic, and other com-	
of fever among labourers	44	plaints of the bowels often	
Posture, injurious effect of,		occasioned by the same	
long inclined - - -	47	cause as - - -	<i>id.</i>
—, figure of body, how		Saint Vitus's dance - - -	313
hurt by - - -	<i>id.</i>	Salsafy, skirrets, and sever-	
Potatoes considered as a		al kinds of beets, observa-	
substitute for bread	30	tions on - - -	33
Potash. <i>See</i> Alkalies.		Sanctorius, an Italian phy-	
Poultices - - -	xxiv	sician quoted on pulmona-	
Powders - - -	<i>id.</i>	ry and cutaneous transpi-	
Powder, astringent - - -	<i>id.</i>	ration - - -	94
—, of bole - - -	<i>id.</i>	Scarborough waters - - -	476
—, carminative - - -	<i>id.</i>	Scald head - - -	428
—, diuretic - - -	<i>id.</i>	Scenes, the variety of, why	
—, aromatic opening,	<i>id.</i>	designed - - -	89
—, saline laxative	xxv	Scirrhus - - -	336
—, steel - - -	<i>id.</i>	Scrofula. <i>See</i> King's Evil.	
—, sudorific - - -	<i>id.</i>	Scurvy - - -	281
—, worm - - -	<i>id.</i>	Sea-sickness - - -	494
—, purging - - -	<i>id.</i>	—, imperial ma-	
—, for tape-worm	<i>id.</i>	rine tincture for - - -	495
Pregnancy, diseases of	403	Sedentary, studious, and la-	
—, during, three		borious occupations, re-	
stages exist - - -	404	marks on - - -	40
Preserves. <i>See</i> Conserves.		—, sedentary studi-	
Prognosis, what - - -	100	ous - - -	50
Q.		—, advice to - - -	<i>id.</i>
Quickenings - - -	404	—, diseases of - - -	<i>id.</i>
—, opinion rela-		Seltzer water - - -	477
tive to - - -	<i>id.</i>	Senses, disorders of - - -	329
Quinsey. <i>See</i> Sore Throat,		Shortsightedness. <i>See</i> My-	
inflammatory.		opia.	
—, malignant - - -	197	Silver, nitrate of, <i>See</i> Min-	
R.		eral Poisons.	
Rattlesnake, bite of - - -	351	Simples, a list of. <i>See</i> Ap-	
Resurrection. <i>See</i> Animation		pendix.	
suspended.		Simplicity, the great rule of	
		diet - - -	75
		Sinapisms. <i>See</i> Cataplasms.	

- Skin, over distention of, in pregnant women - 408
 Smell and taste - 333
 Shoe, the size and figure how regulated - 73-74
 Sleep, want of, and restlessness of pregnant women 407
 —, its benefits - 69
 —, like diet ought to be duly regulated - *id.*
 —, to procure refreshing 70
 —, light suppers procure *id.*
 —, refreshing, when 71
 Small-pox - 168
 Soda. *See* Alkalies.
 Soldiers and sailors, the consequences of their employment - 44
 Sore throat, inflammatory 193
 Spirit of wine, camphorated - xxviii
 Spring-chalybeate in the Isle of Wight - 489
 Squinting. *See* Strabismus.
 Stale bread, its effects upon the humours - 45
 Stone and gravel - 234
 Stomach, cramp of - 316
 —, pain of - 259
 —, inflammation of 210
 —, and bowels, affections of. *See* Costiveness.
 Studious and intense thinkers, advice to - 52
 —, diet of - 54
 Studious occupations, remarks on - 40
 Strabismus - 331
 Strains - 377
 Strangling. *See* Suffocation.
 Study and exercise, the best time for - 53
 Subjects, gloomy ones, not to be inculcated - 91
 Suffocation from various causes - 393
 — of infants - *id.*
 Suppers and breakfasts contrasted - 61
 Surgery - 352
 Swellings, puffy, of the feet, of pregnant women - 406
 Swooning - 318
 Syncope. *See* Fits, Fainting.
See Swooning.
 Syrups - xxv
 T.
 Tap-rooms, highly pernicious 98
 Taste and smell - 333
 Tea, the mischief of, when substituted for solid food 24
 Teething - 431
 Tests. *See* Poisons.
 Thrush. *See* Aphthæ.
 Tinea, capitis. *See* Scald Head.
 Tinctures - xxvi
 Tincture, imperial marine, against sea-sickness, &c. 495
 —, how taken in nervous and other disorders. *id.*
 —, aromatic - xxvi
 —, compound, of the bark - *id.*
 —, volatile, fœtid *id.*
 —, of guaiacum *id.*
 —, of black hellebore, *id.*
 —, astringent - *id.*
 —, of myrrh and aloes *id.*
 —, of opium - xxvii
 —, sacred, or hiera picra - *id.*
 —, compound of senna *id.*
 —, balsam of tolu *id.*
 —, Spanish flies - *id.*
 —, rhubarb - *id.*
 —, tonic - *id.*
 Tooth-ach - 256
 —, of pregnant women - 406
 Touch, disorders of - 335
 Transitions, sudden, &c. 97
 Tunbridge-wells - 470
 —, medicinal properties - *id.*
 —, Pyrmont, and German spa, compared with - 473
 Turner's cerate. *See* Ointment of Calamine.
 Turnips, how considered as food. - 33
 U.
 Ulcers - 362

Ulcers, cure of by roller and compresses, &c. on Mr. Whately's plan - - -	363	Water, tar - - -	<i>id.</i>
—, Baynton, his plan of curing old - - -	365	—, simple distilled - - -	<i>id.</i>
—, Home, Sir Everard, on - - -	367	—, cinnamon - - -	xxx
—, Dr. White, on - - -	<i>id.</i>	—, peppermint - - -	<i>id.</i>
Urinary organs, diseases of. <i>See Diabetes.</i>		—, spearmint - - -	<i>id.</i>
Urine, suppression of - - -	233	—, rose - - -	<i>id.</i>
—, bloody - - -	248	—, spirituous distilled - - -	<i>id.</i>
—, suppression of, in pregnant women - - -	406	—, Jamaica pepper - - -	<i>id.</i>
—, incontinency of, in pregnant women - - -	407	—, spirituous cinnamon - - -	<i>id.</i>
V.		—, necessary to cleanliness - - -	81
Vaccination - - -	493	Wet clothes - - -	95
Vapours, noxious - - -	385	— feet - - -	<i>id.</i>
Vegetable and animal food, difference between - - -	39	Wheys - - -	xxx
—, observations on - - -	56	Whey, alum - - -	<i>id.</i>
Vinegars - - -	xxviii	—, mustard - - -	<i>id.</i>
Vinegar of litharge - - -	<i>id.</i>	—, scorbutic - - -	<i>id.</i>
—, roses - - -	xxix	Whites. <i>See Fluor Albus.</i>	
—, squills - - -	<i>id.</i>	Wind. <i>See Flatulencies.</i>	
Vomiting of infants - - -	228	Wines, - - -	xxxix
—, when from other affections - - -	229—230	Wine, anthelmintic - - -	<i>id.</i>
W.		—, bitter - - -	<i>id.</i>
Waters by infusion - - -	xxix	—, ipecacuanha - - -	<i>id.</i>
Water, lime - - -	<i>id.</i>	—, chalybeate or steel - - -	<i>id.</i>
—, compound - - -	<i>id.</i>	—, stomach - - -	<i>id.</i>
—, sublimate - - -	<i>id.</i>	Womb, inflammation of - - -	417
—, styptic - - -	<i>id.</i>	Women, diseases of - - -	397
		Woody nightshade. <i>See Poisons, Narcotic.</i>	
		Worms - - -	261
		Wounds - - -	357
		Z.	
		Zinc, sulphate of, &c. - - -	
		—, oxyd of. <i>See Mineral Poisons.</i>	



